

Accepted refereed manuscript of:

Costa Castro LF, Tocher DR & Monroig O (2016) Long-chain polyunsaturated fatty acid biosynthesis in chordates: insights into the evolution of Fads and Elovl gene repertoire, *Progress in Lipid Research*, 62, pp. 25-40.

DOI: [10.1016/j.plipres.2016.01.001](https://doi.org/10.1016/j.plipres.2016.01.001)

© 2016, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Long-chain polyunsaturated fatty acid biosynthesis in chordates: insights into the evolution of Fads and Elovl gene repertoire

L. Filipe C. Castro^{1,2}, Douglas R. Tocher³ and Oscar Monroig^{3*}

¹ CIIMAR – Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Porto, Portugal

² Biology Department, FCUP, University of Porto, Portugal

³ Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9 4LA, Scotland, UK

*Corresponding author

Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9 4LA, Scotland, UK

Tel: +44 1786 467892; Fax: +44 1786 472133; E-mail: oscar.monroig@stir.ac.uk

21 **Abbreviations**

- 22 aa, amino acid
- 23 ACP, acyl carrier protein
- 24 ALA, α -linolenic acid (18:3n-3)
- 25 ARA, arachidonic acid (20:4n-6)
- 26 BHT, butylated hydroxytoluene
- 27 cDNA, complementary DNA
- 28 CoA, Coenzyme A
- 29 DHA, docosahexaenoic acid (22:6n-3)
- 30 ELOVL, elongation of very long-chain fatty acid protein
- 31 EPA, eicosapentaenoic acid (20:5n-3)
- 32 ER, endoplasmic reticulum
- 33 FACES, fatty acid chain elongation system
- 34 FADS, fatty acyl desaturase
- 35 FAE1, fatty acid elongase 1
- 36 FAS, fatty acid synthase
- 37 HADC, β -hydroxyacyl-CoA dehydrase
- 38 KAR, β -ketoacyl-CoA reductase
- 39 KCS, β -ketoacyl-CoA synthase
- 40 LA, linoleic acid (18:2n-6)
- 41 LC-PUFA, long-chain (C₂₀₋₂₄) polyunsaturated fatty acids
- 42 ORF, open reading frame
- 43 PKS, polyketide synthase
- 44 PUFA, Polyunsaturated fatty acid
- 45 SCD, stearoyl-CoA desaturase
- 46 TER, trans-2-enoyl-CoA reductase
- 47 VLC-PUFA, very long-chain (>C₂₄) polyunsaturated fatty acid
- 48 WGD, whole genome duplication
- 49

Abstract

Long-chain polyunsaturated fatty acids (LC-PUFA) are major components of complex lipid molecules and are also involved in numerous critical biological processes. Studies conducted mainly in vertebrates have demonstrated that LC-PUFA can be biosynthesized through the concerted action of two sets of enzymes, namely fatty acyl desaturases (Fads) and elongation of very long-chain fatty acid (Elovl) proteins. While LC-PUFA research is a thriving field, mainly focused on human health, an integrated view regarding the evolution of LC-PUFA biosynthetic genetic machinery in chordates is yet to be produced. Particularly important is to understand whether lineage specific life history trajectories, as well as major biological transitions, or particular genomic processes such as genome duplications have impacted the evolution of LC-PUFA biosynthetic pathways. Here we review the gene repertoire of Fads and Elovl in chordate genomes and the diversity of substrate specificities acquired during evolution. We take advantage of the magnitude of genomic and functional data to show that combination duplication processes and functional plasticity have generated a wide diversity of physiological capacities in extant lineages. A clear evolutionary framework is provided, which will be instrumental for the full clarification of functional capacities between the various vertebrate groups.

Keywords

Chordates; elongation of very long-chain fatty acid protein; evolution; front-end desaturase; long-chain polyunsaturated fatty acids

1. Introduction

Unlike proteins and carbohydrates, that generally have structures based on long chains (polymers) of amino acid (polypeptides) or sugar (polysaccharides) residues, lipids are a much more diverse range of compounds with considerable variations in structure. However, although lipids do not have “building blocks” as such, fatty acids come closest, being components of many lipid classes including acylglycerols (glycerides and phosphoglycerides) and sphingolipids. In complex lipids, fatty acids are esterified to alcohol or amino groups and, as these lipid classes are the predominant forms, fatty acids constitute the bulk of lipid. All fatty acids play important roles in key biological processes including energy supply, structure and functions of biological membranes. Some fatty acids, particularly polyunsaturated fatty acids (PUFA) and their derivatives, are highly biologically active and involved in signalling and the regulation of lipid metabolism, inflammatory response and cell division [1]. Saturated and monounsaturated fatty acids can be biosynthesized by all organisms whereas PUFA generally have to be obtained in the diet of animals although they can be converted to long-chain (C₂₀₋₂₄) PUFA (LC-PUFA) in some species. This review will focus on two sets of enzymes, fatty acyl desaturases and elongases that participate in the biosynthesis of LC-PUFA in chordates. In particular, we will review the currently available data on the desaturase and elongase gene¹ repertoire present in chordate genomes and the diversity of substrate specificities that have been acquired by the encoded enzyme proteins during evolution. For clarity purposes, we will first provide a description of the fatty acid nomenclature system used in this paper, as well as a definition of the groups of organisms that compose the chordate phylum.

¹ Gene/protein nomenclature

The standard vertebrate gene symbol formatting determines that different conventions apply to name gene/protein in different model organisms including human (*Homo sapiens*), mouse (*Mus musculus*), rat (*Rattus norvegicus*), chicken (*Gallus gallus*), Carolina anole (*Anolis carolinensis*), frog (*Xenopus laevis* or *X. tropicalis*) and zebrafish (*Danio rerio*). Using as example “Elov15”, the human gene is referred as “ELOVL5” and the predicted protein as “ELOVL5”; for mouse and rat, gene will be named as “*Elov15*”, whereas protein will be “ELOVL5”; for chicken and other birds, gene will be termed as “ELOVL5”, whereas protein will be “ELOVL5”; for anole and other reptiles, gene will be termed as “*elov15*”, whereas protein will be “ELOVL5”; for frog and other amphibians, gene will be named as “*elov15*”, whereas protein will be “Elov15”; similarly, for zebrafish and other fish, gene will be named as “*elov15*”, whereas protein will be “Elov15”. For non-vertebrate organisms and agnathans, we have used the same symbols as described for fish (“*elov15*” for genes and “Elov15” for proteins).

1.1 Nomenclature and structure of fatty acids

A fatty acid is essentially an organic molecule with a carboxylic acid group at the end of an aliphatic chain containing four or more carbons, usually an even number up to 24, although odd-numbered and longer carbon chains are also found (Fig. 1). The aliphatic chain can be “saturated”, where all carbon-carbon linkages are single bonds and all other carbon bonds are taken by hydrogen, or “unsaturated”, where some carbons are linked by double bonds. Several systems have been used historically for fatty acid nomenclature, but the most commonly used and internationally accepted is that defined by the International Union of Pure and Applied Chemistry (IUPAC) in the Compendium of Chemical Terminology (IUPAC, 1997). In the *n*-*x* (or “omega *x*”) system of nomenclature, fatty acids are described by the general formula, C:D*n*-*x*, where C = chain length, D = number of ethylenic/double bonds, and *n*-*x* (or ω*x*) indicates the position of the first double bond relative to the methyl end of the chain. Therefore, in this nomenclature 18:0 represents a saturated fatty acid containing an 18-carbon aliphatic chain with no double bonds, and 18:1*n*-9 (18:1ω9) denotes a monounsaturated fatty acid with an 18-carbon aliphatic chain with a single, *cis* double bond 9 carbons from the methyl group. Polyunsaturated fatty acids (PUFA), any fatty acid containing two or more double bonds that are most commonly separated by methylene (CH₂) groups, are represented as in the following example, 20:5*n*-3 (20:5ω3), which denotes a 20-carbon aliphatic chain containing five double bonds with the first situated three carbons from the methyl group (Fig. 1). However, in the present review, the main alternative nomenclature, the Δ*x* (delta-*x*) system, is equally important as this is the one commonly used for specifying activities of the fatty acyl desaturase (Fads) enzymes studied herein. In this nomenclature the double bonds are numbered from the carboxyl end of the molecule and so 20:5*n*-3 is written as 20:5^{Δ5,8,11,14,17}. Thus, a Fads that introduces an ethylenic (double) bond five carbons from the carboxyl end of the aliphatic chain is described as having Δ5 activity. Additionally, fatty acids are often still described using trivial names, often reflecting their main sources, such as palmitic acid (16:0) and oleic acid (18:1*n*-9) from palm and olive oils, respectively. Semi-systematic names, such as eicosapentaenoic acid (EPA; 20:5*n*-3) and docosahexaenoic acid (DHA; 22:6*n*-3), are more useful as they at least indicate the numbers of carbons (e.g. eicosa-, 20) and double bonds (e.g. pentaenoic, 5). It is important to clarify that, while PUFA applies to any fatty acid with at least two double bonds and generally, with chain lengths from 18 carbons or more, we will use the term long-chain PUFA (LC-PUFA) for fatty acids with an aliphatic chain length from C₂₀ to C₂₄, and two or more double bonds. Similarly, very long-

chain PUFA (VLC-PUFA) refer to PUFA with two or more double bonds, with fatty acyl chains $> C_{24}$.

1.2 Classification of chordates

Metazoans, a classification that refers to animals, are characterized by being multicellular and heterotrophic, possessing epithelial cells and having the ability to produce sperm and eggs. Their overwhelming diversity can be organized into defined groups on the basis of phylogenetic relationships. The first major division separates two clades on the basis of their body plan symmetry: the pre-Bilateria (e.g. sponges, coral and jellyfish) and the Bilateria. Within the latter, two other groups can be distinguished based on early embryonic development features: the protostomes (e.g. molluscs) and the deuterostomes (e.g. sea urchins, mammals). In this review we will focus primarily on the phylum chordates that, together with hemichordates and echinoderms, comprise the deuterostomes. Chordates are subdivided into three sub-phyla, namely cephalochordates, tunicates and vertebrates (Fig. 2). Cephalochordates and tunicates are usually referred to as “invertebrate chordates” as they lack vertebrae (Fig. 2). Chordates share a number of characteristics including a dorsal hollow nerve cord, pharyngeal slits, postanal tail and a prominent axial notochord. Over the years a consensus over the phylogenetic relationships between these three lineages emerged, with tunicates now considered the sister clade of the vertebrates [2]. The transition towards vertebrates entailed a significant number of innovations including the presence of an endoskeleton, elaborated brain and distinct head [3]. A fundamental separation within the vertebrates involves the presence or absence of a jaw, which divides the agnathans (lampreys and hagfishes), which lack a jaw, from gnathostomes (cartilaginous fish, teleosts, amphibians, birds and reptiles and mammals) (Fig. 2).

Comparative studies involving species from key phylogenetic lineages within the chordates have shed light over the specific acquisition (and loss) of numerous developmental, morphological and physiological traits [3-5]. An underscoring aspect of this research is the impact of gene duplication, particularly that arising from whole genome duplications (WGD). It is clear today that WGD events took place in early vertebrate evolution, although the number of events and exact timing remains contentious (Fig. 2) [6-9]. Nevertheless, the increment of gene numbers followed by episodes of sub-functionalizations, neofunctionalizations, and gene loss, have all impacted vertebrate physiology and, with regard to the topic of this review, the number and function of genes encoding key enzymes

involved in the biosynthesis of LC-PUFA. Vertebrates have a surprising diversity of species denoting a wide range of adaptations. They also colonize a variety of habitats from aquatic (fresh and salt) to land and air, which have hinted at impacts on lipid metabolism [10]. In the context of the present review, the availability of full genome sequences in species representatives of the main lineages clearly provides the necessary tools to address the evolutionary and functional aspects of LC-PUFA molecular components (Fig. 2).

2. Biosynthesis of fatty acids

2.1. Biosynthesis of saturated fatty acids

The lipogenic pathways are essentially the same in plants and animals including vertebrates although most studies in animals have been conducted in mammals. The primary pathway of lipogenesis is the biosynthesis of fatty acids, which is catalyzed by two cytosolic enzyme systems, acetyl-CoA carboxylase and the multienzyme fatty acid synthase (FAS) complex that uses acetyl-CoA as carbon source to produce saturated fatty acids, primarily 16:0 in animals and 18:0 in plants [11]. There are two main classes of FAS; Type I systems as found in yeast and animals use a single large, multifunctional polypeptide whereas Type II systems that are found in prokaryotes and plants utilize a series of discrete, monofunctional enzymes. The FAS mechanism consists of sequential decarboxylative Claisen condensation reactions with the key step being the condensation of malonyl-CoA, formed by acetyl-CoA carboxylase, with the growing acyl chain. After each round of elongation the β -keto group is reduced to the fully saturated carbon chain by the sequential action of ketoacyl reductase, dehydrase, and enoyl reductase activities. The growing acyl chain is carried between the enzyme active sites by the acyl carrier protein (ACP) domain through a covalent linkage to the phosphopantetheine prosthetic group, and is finally released as 16:0 by the action of a thioesterase. Acetyl-CoA is initially produced in mitochondria from both carbohydrate or protein sources through the oxidative decarboxylation of pyruvate or the oxidative catabolism of some amino acids, respectively, leading to citrate production (via the tri-carboxylic acid cycle) and export to the cytosol, and subsequent production of cytosolic acetyl-CoA through the action of ATP-citrate-lyase. The NADPH reducing equivalents are produced by enzymes of carbohydrate metabolism including the pentose phosphate pathway (glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase), tricarboxylic acid cycle (NADP-dependent isocitrate dehydrogenase) and malic enzyme.

The FAS complexes are very similar to another family of multi-enzyme complexes, the polyketide synthases (PKS), that use a similar mechanism and homologous domains to produce structurally complex organic molecules or secondary metabolites, collectively termed polyketides, in bacteria, fungi, and plants [12]. Both FAS and PKS have two principal classes, Type I and Type II, with the differences between PKS Types being the same as described above for FAS, e.g. Type II being characterized by discrete, monofunctional enzymes. The relationship between FAS and PKS is particularly interesting and relevant to the present review for two main reasons. Firstly, the evolutionary history of FAS is intimately linked with that of PKS as the Type I FAS in animals is believed to have evolved by modification of fungal Type I PKS, whereas Type I FAS in fungi (and some bacteria) appears to have arisen through the fusion of Type II FAS genes. Secondly, LC-PUFA including EPA and DHA may be biosynthesized by PKS-like pathways (see below) in some marine microalgae including some dinoflagellates and thraustochytrids [13].

2.2 Fatty acid chain elongation

Although FAS produces 16:0 and 18:0, C₂₀ - C₂₄ fatty acids are common and chain lengths > C₂₄ are often found in eukaryotes [14, 15]. Therefore, fatty acid elongation is another major pathway in the production of fatty acids. Depending upon the organism, different terms such as “elongase” or “elongase system”, or fatty acid chain elongation system (FACES) have all been used to describe enzymes responsible for the elongation of fatty acids through the addition of two carbon units to the carboxyl end of the chain. In higher plants, elongation is usually restricted to saturated and monounsaturated fatty acids and accomplished by an enzyme coded by the FAE1 gene (Fatty acid elongase 1) [16], although a PUFA Fae1 has been functionally characterized in the marine parasitic protozoon *Perkinus marinus* [17]. Marine microalgae rich in n-3 LC-PUFA appear to have an elongase (e.g. IgASE1 in *Isochrysis galbana*) capable of elongating PUFA [18]. Similarly, marine protists such as thraustochytrids may also have an elongase capable of elongating PUFA [19]. The yeast, *Saccharomyces cerevisiae*, only produces saturated and monounsaturated fatty acids and has a family of elongases, ELO1, ELO2 and ELO3, with different fatty acid specificities according to chain length [19, 20]. The PUFA-producing fungus, *Mortierella alpina*, that produces arachidonic acid (ARA; 20:4n-6) has two elongases including GLELO that elongates 18:3n-6 to 20:3n-6 and is likely the rate-limiting step in ARA biosynthesis, and MAELO that preferentially elongates saturated and monounsaturated fatty acids [21].

In plants and animals, the elongation of fatty acids to produce fatty acids with chain lengths $> C_{18}$ involves four sequential reactions to add each 2-carbon unit that occurs in endoplasmic reticulum (ER) [19, 22] (Fig. 3). These reactions are analogous to those of *de novo* synthesis, (specifically condensation, 1st reduction, dehydration and 2nd reduction) and are catalyzed by four membrane-bound enzymes, namely Elongation of very long fatty acids (Elovl) proteins (condensing enzyme), β -ketoacyl-CoA reductase (Kar; 1st reduction), β -hydroxyacyl-CoA dehydrase (Hadc; dehydration), and trans-2-enoyl-CoA reductase (Ter; 2nd reduction) (Fig. 3). Early biochemical studies provided indirect evidence to indicate that the condensing enzyme Elovl of the elongation system was rate-limiting, regulated the specificity of elongation in term of chain length and the degree of unsaturation, and was regulated by nutritional and hormone status [19]. Cloning of the condensing enzyme and functional characterization of the activity confirmed these data and other enzymes in the complex do not show any fatty acid specificity [20, 23]. While expanded in Section 3, it is worth clarifying that ELOVL enzymes comprise a family with at least seven members, ELOVL1-7, that can be broadly sub-divided into elongases of saturated and monounsaturated fatty acids, ELOVL1, ELOVL3, ELOVL6 and ELOVL7, and elongases of PUFA, ELOVL2, ELOVL4 and ELOVL5 [20, 22, 24]. It is worth clarifying that Elovl and Fae1 are both β -ketoacyl-CoA synthases (Kcs) and, although structurally unrelated, both enzymes have been suggested to similarly interact with the “core” elongase components Kar, Hadc and Ter (Fig. 3) [25]. Interestingly, a new component of the plant elongation pathway, the ECERIFERUM2-like proteins, are believed to affect the Fae1-like Kcs although the mechanism is not yet been established [26]. Preliminary *in silico* analyses appear to suggest that such a paralogue does not exist in non-plant organisms.

2.3 Biosynthesis of unsaturated fatty acids

The biosynthesis of unsaturated fatty acids can occur by two pathways. The first pathway, found in many bacteria, generates the double bonds by essentially leaving those created during the biosynthesis of the fatty acid [27]. Some marine bacteria (e.g. *Shewanella* sp.) are capable of EPA production and analysis of the amino acid sequences derived from the genes responsible showed they were related to microbial PKS (and FAS) complexes [28]. Related genes have been analysed from DHA-producing organisms, the bacterium *Moritella marina* strain MP-1 [29] and *Schizochytrium*, a thraustochytrid marine protist [30]. The second pathway, found in almost all eukaryotes and some bacteria, involves aerobic desaturase enzymes that directly regioselectively introduce double bonds into fatty acids produced by *de*

novo synthesis or obtained through the diet [31, 32]. Although the basic mechanism of the desaturation reaction, removing two hydrogens to create a carbon-to-carbon double bond, is similar and they all utilize molecular oxygen and reducing equivalents obtained from an electron transport chain, fatty acid desaturases have been classified in a number of ways.

2.3.1 Classification of fatty acid desaturases

Probably the most fundamental classification of fatty acid desaturases is based on subcellular location, which classifies fatty acid desaturases into two groups, the soluble desaturases and membrane-bound desaturases, that are phylogenetically unrelated [31]. The only soluble desaturase is the acyl-acyl carrier protein (ACP) or stearyl-ACP desaturase, found only in the stroma of plant plastids, which is responsible for the production of oleic acid (18:1n-9) from stearic acid (18:0) [33]. In contrast, membrane-bound desaturases are ubiquitous in eukaryotes and bacteria, and are characterized by three histidine box motifs that contain eight histidine residues [34]. The membrane-bound desaturases can be sub-divided based on the nature of the lipid substrate to which the fatty acyl chain is linked [35]. First, acyl-lipid desaturases, which are associated with the ER and chloroplast membrane in plants, cyanobacterial thylakoid membranes, plasma membrane of some bacteria [36], and ER in *Tetrahymena* [37], desaturate fatty acids esterified to glycerolipids. Second, acyl-CoA desaturases, found in the ER of animals and fungi, and plasma membrane of some bacteria, introduce double bonds into fatty acids esterified to coenzyme A (CoA) [38]. A separate classification is based on which end of the fatty acid molecule the desaturase “counts” from in determining specificity. Thus, methyl end desaturases insert the double bond at a fixed number of carbons from the methyl group and are termed ω -desaturases (e.g. ω 3 or ω 6 desaturases), whereas the so-called “front-end” desaturases insert the double bond as a fixed number of carbons from the carboxyl group and are termed Δ (delta)-desaturases (e.g. Δ 6 or Δ 5 desaturases). In Section 3, particular attention will be paid to front-end desaturases termed FADS in mammals. The protein sequences of the FADS-like desaturases possess features characteristic of all membrane-bound fatty acid desaturases, including the three histidine boxes mentioned above, two transmembrane regions, and an N-terminal cytochrome b_5 domain containing the haem-binding motif, HPGG [34, 39, 40]. Thus, the animal front-end desaturases are fusion proteins containing both desaturase and cytochrome b_5 functions [41].

2.3.2 Biosynthesis of monounsaturated fatty acids and C_{18} PUFA

Monounsaturated fatty acids, such as 18:1n-9 and 16:1n-7, are produced through the activity of Δ^9 fatty acyl desaturases, including the soluble stearoyl-ACP desaturase in plants, and the microsomal stearoyl-CoA desaturases of yeast (OLE1) and animals (SCD) [42-44] (Fig. 4). Two representatives of mammalian SCD have been crystallized and subjected recently to X-ray analysis [45, 46]. Therefore, this activity is ubiquitous and found in all living organisms [47]. In plants and algae, monounsaturated fatty acids can be further desaturated through the action of chloroplast membrane-bound desaturases to form, firstly, linoleic acid (LA; 18:2n-6) from 18:1n-9 and then α -linolenic acid (ALA; 18:3n-3) from LA [42, 48] (Fig. 4). The production of LA is catalyzed by an ω^6 desaturase that counts from the pre-existing Δ^9 bond (rather than the methyl end) in higher plants, and a Δ^{12} desaturase in cyanobacteria [43, 49]. The production of ALA from LA is catalyzed by ω^3 or Δ^{15} desaturases [42, 50]. In addition, PUFA can be produced by the action of Δ^{12} desaturases in some yeast [51] and bifunctional desaturases showing both Δ^{12} and Δ^{15} activities have also been described in protozoans [52, 53] and fungi [54].

In contrast to plants, fungi and some aquatic microorganisms, few other organisms are capable of producing PUFA *de novo*. In animals, the Δ^{12} desaturation of 18:1n-9 to form LA was demonstrated in axenic tissues from insects including cockroach [55] and cricket [56], and insect Δ^{12} desaturase genes were subsequently isolated and characterized [57, 58]. The nematode, *Caenorhabditis elegans* has also been shown to have two desaturase-like genes, *fat1* and *fat2*. The *fat1* gene encodes a novel ω^3 desaturase capable of introducing an ω^3 double bond into both C₁₈ and C₂₀ substrates [59], and the *fat2* gene encodes a functional Δ^{12} that can desaturate both C₁₈ and C₁₆ substrates [60], and has been shown recently to also have Δ^{15} desaturation activity [61]. However, PUFA cannot be synthesized *de novo* by chordates/vertebrates as they lack the necessary Δ^{12}/ω^6 and Δ^{15}/ω^3 desaturases [38]. The desaturase enzymes involved in the biosynthesis of monounsaturated and C₁₈ polyunsaturated fatty acids are important in defining context, however the Δ^9 , Δ^{12}/ω^6 and Δ^{15}/ω^3 desaturases are outwith the scope of the present review.

2.4 Biosynthesis of LC-PUFA in plants, bacteria and fungi

De novo biosynthesis of LC-PUFA with chain lengths longer than C₁₈ occurs in phototrophic and heterotrophic algae, especially marine microalgae that are rich in EPA and/or DHA, and some marine bacteria. In algae, LC-PUFA can be produced by two

pathways, either the aerobic pathway using specific, discrete desaturase enzyme proteins [18, 51] or, in some cases including certain thraustochytrids and possibly flagellates, anaerobically via polyketide (PKS) pathways [13]. The aerobic pathway synthesises LC-PUFA from the C₁₈ PUFA produced through the action of the $\Delta 12/\omega 6$ and $\omega 3$ desaturases as described above (Section 2.3). However, the desaturases involved in LC-PUFA biosynthesis are termed “front-end” desaturases as they introduce double bonds into the fatty acyl chains between the $\Delta 9$ bond and the carboxyl end of the molecule, in contrast to the $\Delta 12/\omega 6$ and $\omega 3$ desaturases that produce C₁₈ PUFA, which introduce double bonds between the $\Delta 9$ double bond and the methyl end of the chain [62]. In eukaryotic algae, $\Delta 8$, $\Delta 6$, $\Delta 5$ and $\Delta 4$ desaturases necessary for the production of EPA and DHA from 18:3n-3 have all been described [18] (Fig. 4).

The biosynthesis of n-3 LC-PUFA in marine bacteria also occurs through the activity of a bacterial PKS-type multi-enzyme complex [63]. The PKS pathway for LC-PUFA biosynthesis operates in some thraustochytrids, including *Schizochytrium* sp. [30], and is speculated to also possibly be responsible for DHA production in some flagellate microalgae [13]. In bacteria, the EPA biosynthesis gene cluster carries around 18 open reading frames (ORF), but only five of the genes termed pfaA, pfaB, pfaC pfaD and pfaE code for proteins required for the biosynthesis of EPA [63]. Analysis of the amino acid sequences derived from the genes showed that they were related to microbial PKS (and FAS) complexes [28]. Organization of the gene clusters is divided into three types with Type I, present in *Shewanella pneumatophori* SCRC-2738, containing all five pfa genes in close vicinity. Type II, as in *Moriella marina* MP-1, consists of a cluster of four genes pfaABCD with a separate pfaE, and Type III is considered to consist of just four genes with pfaE integrated into pfaC/E [27].

Biosynthetic enzymes including front-end desaturases and PUFA elongases activities have been also reported in fungi. The filamentous fungus *Mortierella alpina* has an elongase (GLELO) with the ability to elongate 18:3n-6 to 20:3n-6 and thus involved in ARA biosynthesis [21]. Additionally, desaturases with $\Delta 5$ [64] and $\Delta 6$ [65, 66] activities have been also identified. Other fungi from which desaturase genes have been identified include *Rhizopus nigricans*, *R. arrhizus*, *Mucor rouxii* and *Pythium irregulare* [67].

3. Fatty acyl desaturases and their genes in Bilateria

Other than the examples of the nematode *C. elegans* and some insects (Section 2.3.2), bilaterian organisms appear to lack enzymes capable of producing C₁₈ PUFA from monounsaturated fatty acids [39]. Therefore, the production of LC-PUFA in most animals, and certainly in all vertebrates, is restricted to biosynthesis from pre-existing (dietary) C₁₈ PUFA. The pathway for the production of LC-PUFA from C₁₈ PUFA using aerobic front-end fatty acid desaturases (Fads) and fatty acid elongases (Elovl) is shown in Figure 4. The PUFA substrates for the pathway are 18:3n-3 and 18:2n-6 with DHA and 22:5n-6, respectively, generally being the most highly unsaturated products [68]. All the enzymes involved act on both the n-3 and n-6 series fatty acids with generally a preference for n-3 [38]. Using the n-3 example, production of EPA requires $\Delta 6$ and $\Delta 5$ desaturases producing 18:4n-3 from ALA and EPA from 20:4n-3, respectively. An alternative pathway for the production of 20:5n-3 operates via elongation of ALA to 20:3n-3 followed by $\Delta 8$ - and $\Delta 5$ -desaturation steps [69, 70]. In some basal vertebrate lineages, such as teleosts, the production of DHA from EPA can occur directly via a $\Delta 4$ desaturase that produces DHA from the EPA elongation product, 22:5n-3 [71]. In mammals, the production of DHA from EPA appears to involve a more complicated pathway that involves sequential elongations of EPA to produce 24:5n-3 that is desaturated by $\Delta 6$ desaturase [72] (Fig. 4). Recently, the human FADS2 gene product was demonstrated to have the ability for direct $\Delta 4$ desaturation of 22:5n-3 to 22:6n-3 [73].

Bilateria including invertebrate and vertebrate organisms possess genes encoding desaturases and elongases involved in biosynthetic pathways of LC-PUFA, although the exact repertoire and function remain to be clarified. In particular, this section will largely focus on front-end desaturases (Fads) with $\Delta 4$, $\Delta 5$, $\Delta 6$ and $\Delta 8$ activities, and Elovl2-, Elovl4- and Elovl5-like elongases found in chordates. Thus, the current knowledge on gene complement and functionalities of Fads and Elovl involved in LC-PUFA biosynthesis will be revised from basal vertebrates like cartilaginous fish to mammals. For the purpose of providing the adequate evolutionary perspective, the Fads and Elovl genes of non-chordate invertebrates will be also described.

3.1 Fads and Elovl in invertebrates and non-vertebrate chordates

Some invertebrate groups, and particular species among them, are becoming increasingly popular model species for comparative genomics and evolutionary developmental biology.

This is particularly true for species that have been noted for the retention of ancestral features of their genome organization as well as gene content to a much greater degree than seen in more traditional invertebrate model species [74-86]. As a result, ever-increasing genomic data of certain species are becoming available that will surely allow the study of genes encoding PUFA biosynthetic enzymes. While an extensive review of the molecular mechanisms of PUFA biosynthesis in marine invertebrates has been published recently [77], it is still relevant for the present review to highlight the most important findings uncovered in non-vertebrate metazoans, to contextualise the gene complement of Fads and Elovl in chordates, the focus of this review.

The nematode *C. elegans* was the first animal from which front-end desaturases, including $\Delta 6$ [41] and $\Delta 5$ [78, 79] desaturases, were isolated and characterized. More recently, further molecular studies on Fads and Elovl genes have been performed on another group of invertebrates, the molluscs. A complementary DNA (cDNA) encoding a Fads enzyme was first isolated from the common octopus *Octopus vulgaris* and shown to have $\Delta 5$ desaturation activity towards saturated and polyunsaturated fatty acyl substrates [80]. Likewise, a full-length cDNA sequence encoding an Elovl-like protein identified from *O. vulgaris* had high homology to vertebrate Elovl2 and Elovl5 enzymes [81]. Consistent with vertebrate Elovl2 and Elovl5 that have demonstrated roles in PUFA biosynthesis in vertebrates [20], the octopus Elovl was also efficient in the elongation of C18 and C20 PUFA substrates. Further investigations on other mollusc class representatives including *Haliotis discus hannai* (Gastropoda) [82] and *Chlamys nobilis* (Bivalvia) [83, 86], have confirmed the presence of Fads and Elovl with similar substrate specificities to those described previously for the *O. vulgaris* genes. More recently, two Fads- and Elovl-like cDNAs have been isolated and functionally characterized from the common cuttlefish *Sepia officinalis* [85]. These results confirm that the Fads and Elovl repertoire in protostome metazoans appear to have conserved functionalities.

Through the interrogation of transcriptome and genome sequences, Surm and co-workers [86] have recently screened a representative list of Fads and Elovl genes in a number of commercially important species within phylum Mollusca. It was confirmed that orthologues of Fads and Elovl were found in all investigated species covering three major classes including gastropods, bivalves and cephalopods. Regarding the Fads complement, cephalopods were found to possess one single Fads gene as previously predicted [80, 85], whereas a lineage-specific gene duplication event accounted for the presence of more than one Fads copy in some of the gastropod and bivalve species investigated. Interestingly, one

group of sequences (Clade A) included all the mollusc Fads that have been functionally characterized to date as $\Delta 5$ desaturases, but none of the Fads-like enzymes from Clade B have yet been functionally characterized [86]. Similar to Fads, gene duplication has played a major role in the evolution of the Elovl gene family in molluscs. Genomic data support the idea that a duplication event in the Elovl gene family likely occurred during the early diversification of bilateria as all three classes considered (cephalopods, bivalves and gastropods) possess sequences orthologous to the Elovl2/5- and Elovl4-like types found in present day vertebrates [86].

The characterization of *elovl/fads* gene repertoire and function in invertebrate deuterostomes including species from phyla such as the hemichordates, echinoderms, cephalochordates and tunicates is poorly detailed (Fig. 5). Here we focus on basal invertebrate chordate lineages. To date, a single *elovl* gene representative has been characterized in the European amphioxus (*Branchiostoma lanceolatum*) [87] and the sea squirt (*Ciona intestinalis*) [88]. Despite their clear orthology towards the larger super-family of ELOVL enzymes, phylogenetic analysis indicated that these characterized sequences belonged to different Elovl classes. Independent phylogenetic reconstructions demonstrated that the isolated sequence from the sea squirt belonged to the Elovl4 clade [87, 88], while the amphioxus gene was a *bona-fide* orthologue of Elovl2/5 from vertebrates [87]. Perhaps more interesting was the fact that close inspection of the available genomes of the Florida amphioxus (*Branchiostoma floridae*) and two sea squirt species (*C. intestinalis* and *C. savignyi*) suggested a fundamental difference. While orthologues of *elovl4* and *elovl2/5* were recognizable in the lancelet (amphioxus) genome (as they are indeed in protostomes), no apparent *elovl2/5* was found in these tunicate species [87] (Fig. 5). Some data on the characterization of the substrate specificities of these Elovl are also available. The Elovl4 from the sea squirt was able to elongate C₁₈ and C₂₀ PUFA [88]. In contrast, the amphioxus Elovl2/5 showed a more broad elongation pattern with the ability to elongate C₂₂ PUFA, but with clear preferences for C₁₈ and C₂₀ [87]. At present no characterization of the orthologue of Elovl4 has been performed in amphioxus. Regarding desaturases, both amphioxus and sea squirts have a single desaturase gene although their precise phylogenetic relationships and functions have not yet been addressed (Fig. 5) [87, 89]. Preliminary analysis suggests that the tunicate desaturase is not a Fads but rather a sphingolipid desaturase (Castro and Monroig, personal communication).

3.2 Fads and Elovl in vertebrate basal lineages

3.2.1 Agnathans, chondrichthyans and holosteans

The Elovl and Fads gene repertoire in basal vertebrate lineages, such as chondrietyans and especially agnathans, is poorly investigated. Therefore, only patchy knowledge of the functions and diversity of LC-PUFA biosynthetic enzymes outside tetrapods, such as birds and mammals, and teleosts is currently available [90-92]. However, recent findings have provided some clarification of Fads and Elovl genes in early vertebrate evolution, and their overall impact on our understanding of the evolution of the LC-PUFA pathway. The gene complement of *fads* desaturases in chondrietyans was shown to partially coincide with that of mammals [89]. Thus, orthologues of both *fads1* and *fads2* were identified in the elasmobranch, the lesser-spotted dogfish or small-spotted catshark (*Scyliorhinus canicula*), pushing back the origin of these desaturase gene lineages to the pre-gnathostome period [89]. Moreover, the overall PUFA substrate specificity of both gene orthologues showed unequivocally that Fads1 and Fads2 were $\Delta 5$ and $\Delta 6$ desaturases, respectively, similar to that previously determined in mammals [89]. The recent release of the genome sequence of the holocephali species, the elephant shark (*Callorhinchus milii*) [93], confirmed the initial description in the dogfish, with only two Fads-encoding genes found in the chimaera genome (Fig. 6). *Fads* genes tend to occur in genomes in physical clusters, an indication of their origin through tandem gene duplication. That is the case in humans with a gene cluster with three Fads genes, as well as birds and reptiles with four and five Fads genes, respectively [89]. Although the original description of the *fads* repertoire in cartilaginous fish did not provide information regarding the genomic organization of these genes, the *fads1* and *fads2* orthologues in *C. milii* map together into the same genomic scaffold with a similar composition to the human *FADS* gene locus at chromosome 11 (Fig. 6). The finding of *fads1* and *fads2* orthologues in cartilaginous fish, together with the description of a single basal gene in invertebrate chordates [89], provided tentative evidence that both genes originate from a gene duplication dated back to the transition from cephalochordates/tunicates to gnathostomes. In the most basal vertebrate branch, the agnathans, the repertoire of Fads genes is presently unknown (Fig. 5). While the genome of the sea lamprey (*Petromyzon marinus*) presented no detectable gene sequence with similarity to front-end desaturases, the existence of a partial sequence with similarity to *fads2* was suggested previously [89]. Moreover, the recently available genome from the Japanese

lamprey (*Lecentheron japonicus*) [8] indicated the presence of at least a single sequence with similarity to *fads1* (Castro's personal communication) (Fig. 5) and thus the currently available data enable us to tentatively infer the presence of *fads1* and *fads2* orthologues in agnathans. This would tentatively date the origin of these two gene lineages to the time of the invertebrate/vertebrate transition.

The *elovl* gene repertoire involved in LC-PUFA biosynthesis, namely Elov12 and Elov15, has been characterized in agnathans and chondriactyans. Monroig and co-workers [87] have investigated the *elovl2* and *elovl5* gene repertoire and function in two species belonging to these chordate lineages, the sea lamprey and the elephant shark (Fig. 5). In the case of the elephant shark, orthologues of the tetrapod Elov12 and Elov15 were identified and their origin delineated using both phylogenetics and comparative genomics [87]. However, agnathan sequences are often difficult to place phylogenetically [94]. In the case of Elov12/5, supporting evidence from both phylogenetic and synteny comparisons indicated that orthologues of both Elov12 and Elov15 can indeed be found in lampreys [87] (Fig. 5). Analysis of the PUFA substrate specificities of these enzymes revealed a complex evolutionary pattern. The elephant shark Elov1 resembled other gnathostome Elov12 and Elov15 with clear preferences for C₂₂ and C₁₈, respectively, while a clear overlap was observed for C₂₀ PUFA substrates [87]. The profile determined for the lamprey genes was strikingly different. The agnathan putative Elov12 and Elov15 enzymes showed overlapping preferences for C₁₈ and C₂₀, respectively, with no capacity to elongate C₂₂ PUFA [87]. These results suggested that the agnathans represent an intermediate state after duplication since they completely lack the ability to elongate C₂₂ PUFA observed in cartilaginous fish such as elephant shark.

The exact role and gene number of *elovl4* genes in lampreys and cartilaginous fish has not been described or characterized previously. However, genome mining clearly indicated the presence of single copy orthologues in both lineages, which await functional characterization (Fig. 5). Finally, other emerging genomic fish models will most likely provide a better-defined picture of Fads and Elov1 evolution. For example, the impact of the teleost-specific whole genome duplication (3R) has been barely explored considering other non-classical vertebrate models such as the spotted gar, *Lepisosteus oculatus*, a pre-3R holostean [4]. Scrutiny of the spotted gar genome revealed the identification of the full gene complement of Fads (*fads1* and *fads2*) and Elov1 (*elovl2*, *elovl4*, and *elovl5*), an indication that a conserved LC-PUFA pathway is probably present (Castro's personal communication).

3.2.2 Teleosts

The mechanisms of LC-PUFA biosynthesis in teleost fish, particularly farmed species, have been investigated extensively in recent years, and many aspects of these metabolic pathways are better understood in fish compared to mammals. These investigations have led to the conclusion that LC-PUFA biosynthetic capability varied notably among species linked to differences in the complement of Fads and Elovl gene and enzymatic activities existing in that species [92]. Generally, freshwater and salmonid species have been regarded as having high capacity for LC-PUFA biosynthesis, whereas marine teleost species showed a limited ability to biosynthesize LC-PUFA [95]. This has been hypothesized to be a consequence of marine species having evolved in an LC-PUFA-rich environment and thus with low evolutionary pressure to retain the ability to endogenously produce LC-PUFA. In contrast, lower levels of LC-PUFA in the food chain may have been the selective pressure driving freshwater species to retain the ability to biosynthesize LC-PUFA to satisfy their physiological requirements [89, 95, 96]. In addition to the fact that such dichotomy had been established based on a rather limited number of species, other confounding factors beyond habitat have been explored in recent years. Among them, the trophic level, the position of an organism within the food web, the “trophic ecology” and diadromy have been also suggested as potential drivers modulating the range of desaturation and elongation capabilities in teleosts and thus their ability for LC-PUFA biosynthesis [97, 98]. Importantly for the purpose of this review, it has become obvious that the capability for LC-PUFA biosynthesis in teleost fish is more diverse than in other vertebrate groups, and is possibly the result of a combination of factors that interact throughout the evolutionary history of each particular group or species.

3.2.2.1 Fads in teleosts

Teleosts have been by far the group of tetrapods where the greatest number of Fads and Elovl have been studied (Fig. 5). The first Fads-encoding cDNAs were isolated from rainbow trout [99] and zebrafish [100]. The amino acid sequence homology of the two Fads indicated they both were $\Delta 6$ desaturases. However, the functional characterization of the zebrafish Fads by heterologous expression in yeast demonstrated that the encoded enzyme was a bifunctional $\Delta 6/\Delta 5$ desaturase [100]. Subsequently, a relatively large number of Fads-like cDNAs have been isolated from a variety of teleost fish from a wide range of environments (from

freshwater species such as zebrafish, to diadromous species like Atlantic salmon, and marine species such as Atlantic cod), trophic levels (from true herbivores like rabbitfish to top carnivores like Atlantic bluefin tuna) and phylogenetic groups (from the basal teleost clade Elopomorpha such as eels, to Percomorpharia like cobia or gilthead seabream). An extensive list of all Fads-like cDNAs that have been isolated and functionally characterized from teleost fish species is shown in Table 1 [70, 71, 92, 97-117].

It has been shown that all the Fads-like desaturases characterized in teleost fish are *FADS2* orthologues [89]. Interesting, whereas mammalian *FADS2* are predominantly $\Delta 6$ desaturases, teleostei Fads2 enzymes show much more varied substrate specificities (Table 1). Similar to mammalian orthologues, many fish Fads2 are indeed typical $\Delta 6$ desaturases as demonstrated in a range of marine (gilthead sea bream, turbot, Atlantic cod, cobia, European sea bass, barramundi, Northern bluefin tuna, meagre, Nibe croaker) and freshwater species (common carp and rainbow trout). None of those Fads2 showed $\Delta 5$ or $\Delta 4$ desaturase activities. In Atlantic salmon, however, four distinct genes encoding Fads2 have arisen as a result of segmental duplications and not as a result of the salmonid tetraploidization event hypothesized to have occurred between 25 and 100 million years ago [104]. While the so-called $\Delta 6$ Fad_a, $\Delta 6$ Fad_b, and $\Delta 6$ Fad_c [104] were shown to have $\Delta 6$ specificity [103, 104], a fourth Fads2 cDNA was demonstrated to have $\Delta 5$ -desaturase activity [102]. The salmon $\Delta 5$ Fads2, together with the three $\Delta 6$ -like desaturases described above, enable this species to perform all desaturation reactions (i.e. $\Delta 6$ and $\Delta 5$ desaturations) required in the LC-PUFA biosynthesis through the conventional pathways for mammals (Fig. 4). Thus, biosynthesis of DHA in Atlantic salmon appears to proceed through the Sprecher pathway. This is also the case for rainbow trout, for which another monofunctional $\Delta 5$ Fads2 has been characterized (Hamid's personal communication), and zebrafish, the only fish species where there is supporting evidence for the existence of the Sprecher pathway in fish [118-120]. Interestingly, other teleost Fads2 have apparently lost the ability to desaturate $24:5n-3$ to $24:6n-3$, a key step in the Sprecher pathway. Thus, the Nibe croaker *Nibea mitsukurii*, a member of the Sciaenidae family (Percomorpharia) [121], possess a Fads2 that has $\Delta 6$ desaturase activity towards $18:3n-3$ but no capability to desaturate $24:5n-3$ [112]. Whether the apparent loss of $\Delta 6$ desaturase activity on C_{24} substrates in Fads2 of *N. mitsukurii* is an extended trait or not among fish Fads2 desaturases remains to be elucidated. However, it has become apparent in recent years that Fads2 from teleosts have evolved quite markedly from the typical Fads2 functionalities as described in mammals as indicated by their high plasticity in terms of substrate specificities. Other than the abovementioned zebrafish bifunctional $\Delta 6\Delta 5$ Fads, and

the salmon and trout $\Delta 5$ Fads2, other subfunctionalized Fads2 have been subsequently characterized in several fish species.

Arguably the most important and interesting finding in our investigations of teleost LC-PUFA biosynthesis in recent years has been the discovery of a Fads2 in rabbitfish (*Siganus canaliculatus*) with capability for $\Delta 4$ desaturation [71], an activity originally believed to account for the production of DHA directly from 22:5n-3 resulting from a single elongation of EPA [68]. More recently Fads2 with $\Delta 4$ activities have also been characterized in Senegalese sole *Solea senegalensis* [97], pike silverside *Chirostoma estor* [92] and the striped snakehead *Channa striata* [117]. Further to the isolation and characterization of $\Delta 4$ desaturase cDNAs in teleosts, *in vivo* biochemical evidence of the existence of an enzymatic activity enabling $\Delta 4$ -desaturase-dependent DHA synthesis has been shown recently in *S. senegalensis* [122]. In comparison with the Sprecher pathway, the so-called “ $\Delta 4$ pathway” is a more direct metabolic route as it avoids translocation of LC-PUFA intermediates (namely 24:6n-3) between endoplasmic reticulum and peroxisomes, and also the further catabolic step (partial oxidation to DHA) occurring in the latter organelle [72, 123]. In addition to the $\Delta 4$ Fads2, rabbitfish also expressed a bifunctional $\Delta 6\Delta 5$ Fads2 [71], similar to the zebrafish desaturase, and the high degree of amino acid (aa) sequence similarity between the two rabbitfish Fads2 has prompted investigation of the specific residues dictating the different substrate specificities of Fads2 desaturases [124]. It was concluded that four aa residues (YNYN for $\Delta 4$ Fads2 and FHYQ for $\Delta 6\Delta 5$ Fads2) located between the second and third histidine boxes were responsible for the PUFA substrate specificities of each enzyme. *In silico* searches among teleost fish indicated that Fads2 with the key aa residues and thus being putative $\Delta 4$ desaturases were more widely spread than initially believed. For instance, cichlids including *Oreochromis niloticus*, *Maylandia zebra*, *Haplochromis burtoni* and *Pundamilia nyererei*, all possess putative $\Delta 4$ Fads2 (Monroig’s personal communication).

As mentioned above, plasticity of teleost Fads2 is not restricted to the existence of monofunctional $\Delta 6$, $\Delta 5$ or $\Delta 4$ desaturases, but some of the enzymes investigated have been shown to be bifunctional proteins. In addition to the abovementioned zebrafish Fads2 that was the first bifunctional $\Delta 6\Delta 5$ desaturase described [100], other bifunctional desaturases were described in fungi [54, 125], protozoans [53] and moths [126]. In fish, the rabbitfish *S. canaliculatus* [71] as described above, Nile tilapia *O. niloticus* and pike silverside *C. estor* [92] also possess bifunctional $\Delta 6\Delta 5$ Fads2 enzymes. These bifunctional desaturases are enzymes that can introduce two separate, distinct double bonds into the same fatty acid chain and, interestingly, mammalian Fads2, zebrafish $\Delta 6\Delta 5$ Fads2 and possibly salmon Fads2_a,

are also believed to be responsible for introducing two distinct bonds into both C₁₈ and C₂₄ chains as required for DHA synthesis [120, 127, 128]. Many Fads2 desaturases are also able to introduce the same double bond into both C₁₈ and C₂₀ chains. Specifically, a retrospective study investigating the ability of Fads2 from freshwater, diadromous and marine teleost species to desaturate PUFA at position Δ8, an intrinsic characteristic of mammalian FADS2 [69], concluded that is also a characteristic of teleost Fads2. That was also the case of bifunctional Δ6Δ5 desaturases, and so this sort of multifunctionality appears to be an extended feature among teleost Fads2. Furthermore, the Δ8 activity varied notably among the species investigated, with marine fish Fads2 having higher Δ8 capability than Fads2 from freshwater/diadromous species. The establishment of Δ8 activity in Fads2 enzymes opened up the possibility of a “Δ8 pathway” whereby 20:3n-3 was no longer regarded as a dead-end product but could potentially be an intermediate in the LC-PUFA biosynthetic pathway. However, it remained unclear what benefit this evolutionary adaptation implied for marine species that are receiving adequate dietary EPA and DHA, and little 18:3n-3 or 20:3n-3, and in which the limiting step in the LC-PUFA biosynthesis pathway appears to be lack of Δ5-desaturation capability [13, 129]. It is important to note that, while the desaturase gene numbers and substrate specificities vary among species, some teleosts appear to lack any *fads*-like gene in their genomes. This is the case of two pufferfish species, namely *Takifugu rubripes* and *Tetraodon nigroviridis* [129].

3.2.2.2 Elovl in Teleosts

Elovl-like cDNAs have been isolated and characterized from a wide range of fish species (Table 2) [92, 96-98, 107, 110-113, 115, 130-140]. The first teleost Elovl-like cDNAs with a role in LC-PUFA biosynthesis was a zebrafish elongase that was identified as an Elovl5 enzyme [130]. Similar to mammalian orthologues [141, 142], zebrafish Elovl5 showed the ability to elongate C₁₈ and C₂₀ PUFA, with markedly lower elongation efficiency on C₂₂ PUFA substrates. Subsequently, Elovl5 cDNAs were cloned and characterized in Atlantic salmon, the only teleost species where two paralogues have been cloned and functionally characterized [102, 134], catfish, tilapia, turbot, gilthead sea bream, Atlantic cod [133], cobia [107], barramundi [110], Southern [137] and Northern [113] bluefin tuna, rabbitfish [138], black seabream [139], meagre [98], rainbow trout [140] and Nibe croaker [112]. While the majority of these studies on the functional characterization of Elovl5 included only 18:4n-3 and 18:3n-6 as substrates for C₁₈ elongation, the more recent studies increasingly include

both 18:3n-3 and 18:2n-6 as potential substrates for elongation [98, 137]. Consequently, Elovl5 has been demonstrated to have the ability to elongate 18:3n-3 and 18:2n-6 and thus produce adequate Δ 8-desaturation substrates 20:3n-3 and 20:2n-6 (Fig. 4). The establishment of the ability of Elovl5 to elongate 18:3n-3 and 18:2n-6 along with the aforementioned Δ 8 activity in Fads2 enzymes has confirmed the possibility of a “ Δ 8 pathway” and so the initial steps in LC-PUFA biosynthesis can be performed by the same Fads2 and Elovl5 enzymes but with either activity initiating the pathway, desaturase followed by the elongase (Δ 6) or elongase followed by desaturase (Δ 8). Interesting, the meagre *A. regius* and the Nibe croaker *N. mitsukurii* Elovl5 have also shown the ability to elongate the C₁₆ PUFA 16:3n-3 [98, 112], a fatty acid found in algae and plants. Moreover, heterologous expression in yeast suggested a putative role for Elovl5 in fish in the elongation of monounsaturated fatty acids, as denoted by the endogenous 18:1n-7 and 18:1n-9 in yeast being elongated to 20:1n-9 and 20:1n-7 [98, 102, 110, 112, 134]. The teleost Elovl5 functionalities suggest that these are more versatile and adaptive enzymes than the mammalian orthologues. In this regard, adaptation to diet was postulated as the main evolutionary driver accounting for the different tissue expression of the two Elovl5 paralogues found in Atlantic salmon, and that of Northern pike (*Esox lucius*), the closest extant relative to the preduplicated ancestral salmonid [96]. Thus, the complementary expression pattern of *elovl5* genes in Atlantic salmon (liver and intestine) might have enabled Atlantic salmon to thrive in a relatively poor LC-PUFA environment during the freshwater stages. It was postulated that Atlantic salmon *elovl5* genes have been efficiently retained in the genome under strong functional constraints indicating a physiological requirement for both enzymes to be functionally active. On the other hand, a piscivorous species like the Northern pike, despite it has the ability to biosynthesize DHA from ALA [143], can directly satisfy its LC-PUFA requirements in the diet and thus the Elovl5 activity for LC-PUFA production in liver and intestine might not be so critical. Upstream promoter regions of salmon *elovl5* have strongly diverged from one another, and thus, in contrast to their similar substrate preferences, the different regulatory mechanisms indicate a relaxation of purifying selection following the duplication event [144].

Investigations of the other Elovl enzymes involved in LC-PUFA biosynthesis, i.e. Elovl2 and Elovl4, has contributed considerably to an advanced understanding of these metabolic pathways in teleosts (Fig. 5). An Elovl2 enzyme was cloned and functionally characterized for the first time in Atlantic salmon [134] and soon thereafter in zebrafish [131]. While showing some low level of elongation activity towards C₁₈ substrates (18:4n-3 and 18:3n-6), the Elovl2 were particularly efficient in the elongation of C₂₀ (20:5n-3 and 20:4n-6) and C₂₂

(20:5n-3 and 20:4n-6) LC-PUFA substrates, the latter not regarded as substrates for Elovl5 according to the low (if any) conversions observed in the yeast expression system. Recently, an Elovl2 has been isolated and investigated in rainbow trout *O. mykiss* [140]. Similar to the observations in *S. salar* and *D. rerio*, the *O. mykiss* Elovl2 showed the ability to elongate C₂₀ and C₂₂ but, in contrast, did not show any activity on C₁₈ PUFA substrates. Interestingly it is unclear whether the ability to elongate C₁₈ PUFA is a common feature among Elovl2 proteins as, like Atlantic salmon and zebrafish orthologues, the mouse ELOVL2 showed some elongation capability on C₁₈ PUFA [141], although the human and chicken did not (see below), similar to rainbow trout Elovl2 [90, 142]. Regardless of its possible role in C₁₈ PUFA elongation, the ability of Elovl2 to elongate 22:5n-3 to 24:5n-3 was regarded as key for the production of DHA via the “Sprecher pathway”, and because of the loss of Elovl2 in the vast majority of marine fish lineages and virtually all commercially important farmed species, this has been hypothesized as a possible factor accounting for the limited ability for DHA biosynthesis in those species [134].

Elovl4 are the most recent member of the Elovl family that has been studied in teleosts. Zebrafish was the first teleost fish species from which Elovl4 were isolated and functionally characterized, representing at that time the only report of a non-human Elovl4 elongase [132]. Two isoforms, termed as Elovl4a and Elovl4b, were described in zebrafish and both showed the ability to efficiently elongate saturated fatty acids up to C₃₆. Interestingly though, activity for the elongation of PUFA substrates was only shown by Elovl4b, which effectively converted the C₂₀ LC-PUFA, EPA and ARA, to elongated products with acyl chain-lengths up to C₃₆. Subsequent studies describing the cloning and functional characterisation of Elovl4 were conducted in cobia (*R. canadum*), Atlantic salmon (*S. salar*) and rabbitfish (*S. canaliculatus*) [135, 136, 138]. In agreement with these enzymes being homologues of the zebrafish Elovl4b isoform, the cobia, Atlantic salmon and rabbitfish Elovl4 enzymes showed the ability to biosynthesize both saturated and polyunsaturated VLC-FA. While Atlantic salmon also possess an Elovl2 with elongation capability towards C₂₂ LC-PUFA [134], the ability of Elovl4 from cobia and rabbitfish, two marine Actinopterygii species, to elongate 22:5n-3 to 24:5n-3 provided evidence to support a role for Elovl4 of some teleosts at least in DHA biosynthesis similar to Elovl2 [136]. More recently, the Nibe croaker Elovl4 [112], another Elovl4b-like orthologue, was shown to be capable of elongating 22:5n-3 to 24:5n-3 although the reported conversions were notably lower (3.6 %) compared to previously reported conversions for zebrafish isoform b (29.8 %), cobia (34.1 %), Atlantic salmon (22.1 %) and rabbitfish (20.7 %). However, these studies on fish Elovl4 elongases have

revealed that Elovl4 from some marine species can elongate 22:5n-3 to 24:5n-3 and other C₂₂ LC-PUFA, and this has been hypothesized as an adaptive strategy to compensate for the lack of Elovl2 in these species [136]. *In silico* searches strongly suggest that all teleosts appear to possess at least one copy of both *elovl4a* and *elovl4b*. It is uncertain which mechanism has led teleosts to have two distinct genes for Elovl4 enzymes, but it is clear that they may have diverged to acquire different functions and tissue distributions [132]. A comprehensive tissue distribution study on ten *elovl* genes performed in Atlantic cod (*Gadus morhua*) recently showed that, in addition to Elovl4a and Elovl4b homologues, this teleost possessed two further *elovl4*-like transcripts termed as *elovl4c-1* and *elovl4c-2* [145]. Further studies investigating the presence of this Elovl in other teleost species and its function are required.

3.3 Fads and Elovl in non-mammalian tetrapods

This section reviews the existing knowledge on LC-PUFA biosynthetic enzymes in reptiles, birds, and amphibians. Basal tetrapods such as amphibians display the complete repertoire of Fads and Elovl although their function has not been elucidated (Fig. 5). In birds and reptiles an interesting finding was described with respect to the Fads gene collection as investigation into the genomes of representative species showed a unique Fads1 gene expansion in this lineage (Fig. 5) [89]. It is unclear at this stage whether these enzymes perform different functions, for example utilizing alternative PUFA substrates (sub-functionalization), or if they have evolved novel functions with respect to PUFA desaturation. Since this gene expansion dates back to reptile/bird separation we anticipate that these genes will perform similar functions in both lineages. Additionally, given the relevance of poultry production it is important that future studies investigate Fads capacities in chicken.

The chicken (*Gallus gallus*) ELOVL2 and ELOVL5 were the first elongase genes to be studied in birds [90] (Table 3). Functional characterization of the two proteins in the heterologous yeast *S. cerevisiae* showed the ELOVL2 was capable of elongating C₂₀-C₂₂ PUFA substrates, and no activity on C₁₈ PUFA such as 18:4n-3 and 18:3n-6 being detected. In contrast, the functionality of the chicken ELOVL5 was reported to be broader compared to ELOVL2. Thus, the chicken ELOVL5 showed the ability to elongate a variety of substrates including C₁₈ (18:4n-3 and 18:3n-6), C₂₀ (EPA and ARA) and, interestingly, C₂₂ (22:5n-3 and 22:4n-6) PUFA. Indeed, the ability of the chicken to produce 24:5n-3 from EPA was regarded as unique when compared to non-piscine orthologues such as human [142] and rat [146]. However, as discussed above (Section 3.2.2.2), Elovl5 from some teleost fish showed some

limited ability to elongate C₂₂ PUFA including 22:5n-3 and, as suggested for the chicken ELOVL5, they can potentially contribute to the biosynthesis of DHA through the Sprecher pathway [72].

The ability of the chicken ELOVL5 to elongate C₂₂ PUFA does not appear to be a common feature among bird ELOVL5. Thus, a subsequent study on fatty acid elongases of the domestic Mallard duck (*Anas platyrhynchos*) and Australian white hybrid turkey (*Meleagris gallopavo*) showed that the Elov15 from these species had no activity towards C₂₂ PUFA including 22:5n-3 and 22:4n-6 [91]. Unlike the chicken ELOVL5, both the duck and turkey ELOVL5 had C₁₈ PUFA such as 18:4n-3 and 18:3n-6, as well as C₂₀ PUFA like EPA and ARA as preferred substrates for elongation. Similarly, functional analyses of the duck and turkey ELOVL2 also showed some differences compared to the chicken orthologue [90]. Although the turkey ELOVL2 did not show any activity towards C₁₈ PUFA substrates like the chicken ELOVL2, the turkey orthologue also showed no activity on ARA that remained unmodified. The duck ELOVL2 was capable of elongating C₂₀ (including EPA and ARA) and C₂₂ substrates similar to the chicken ELOVL2 but, in addition, it showed the ability to elongate C₁₈ PUFA like 18:4n-3 and 18:3n-6, an capability not shown by the chicken orthologue. The ability of Elov12 proteins to elongate C₁₈ PUFA substrates to a small extent has also been described in fish as detailed above [131, 134] and mouse [142].

3.4. FADS and ELOVL in mammals

The LC-PUFA biosynthetic enzymes have been extensively studied in human and other mammals, primarily rodent model species such as mouse *Mus musculus* and rat *Rattus norvegicus*. As a result, several reviews have been published in recent years [20, 22, 32]. Therefore, this section will provide a brief overview of the *FADS* and *ELOVL* gene complement in mammals and the functions of their protein products with a role in LC-PUFA biosynthesis. Additionally, we will also discuss the most recent advances on evolutionary aspects of these gene families.

3.4.1 Mammalian *FADS*

Mammals express $\Delta 6$ and $\Delta 5$ desaturases with the ability to desaturate at the $\Delta 6$ and $\Delta 5$ positions of PUFA substrates. In humans the $\Delta 5$ and $\Delta 6$ desaturase enzymes are encoded by the *FADS1* and *FADS2* genes, respectively [147] (Fig. 5). These genes are organized into a

tight physical cluster on chromosome 11, and a further desaturase, FADS3 with unknown function [148], is also part of the cluster [147, 148]. Interestingly, Castro and co-workers [89] also identified a fourth Fads gene (*Fads4*) sequence occurring at a distant genomic location of the *FADS1/FADS2/FADS3* gene cluster in mouse. The function and role of murine *Fads4* remains to be characterized, but its unique phylogenetic distribution and rather restricted occurrence in some mammalian species, perhaps suggests a very specific role.

By comparison of the sequences of known $\Delta 6$ desaturases from a cyanobacterium [150], nematode [41] and plant [151], full lengths cDNAs for $\Delta 6$ FADS2 desaturases were characterized in mammals including rat [152], mouse and human [153] (Table 3). Functional characterization demonstrated that the C_{18} PUFA, 18:3n-3 and 18:2n-6, were adequate substrates for the mammalian $\Delta 6$ desaturases and, subsequently, it was demonstrated that the $\Delta 6$ desaturase from human [127] and rat [128] also had the ability to desaturate 24:5n-3 at the $\Delta 6$ position [127]. While the recent discovery of $\Delta 4$ activity of the human FADS2 provides an alternative and direct pathway for DHA biosynthesis [73], historically it was accepted that DHA production was achieved through the Sprecher pathway and, thus, FADS2 mediates the $\Delta 6$ desaturation of 24:5n-3 to 24:6n-3, which is thereafter translocated to peroxisomes for chain shortening via partial β -oxidation [123]. This makes DHA biosynthesis by the Sprecher pathway a somewhat inefficient process and thus $\Delta 6$ desaturase has been regarded as the overall rate-limiting factor in the LC-PUFA biosynthetic pathway in mammals [22]. Unlike $\Delta 6$ desaturases, the $\Delta 5$ desaturases operate at one single step of the pathway, converting 20:4n-3 to EPA in the n-3 series, and 20:3n-6 to ARA in the n-6 series (Fig. 4). Several mammalian $\Delta 5$ FADS1 desaturase genes have been characterized from human [154], rat [155] and mouse [156]. Interestingly, non-coding RNA genes, which are transcribed from the opposite strand of the mammalian *FADS1* gene, were later discovered in human, mouse and rat [157]. Such reverse $\Delta 5$ -desaturase was found to regulate the expression and enzymatic activity of the rat FADS1, supporting a significant role as a natural antisense regulator of $\Delta 5$ -desaturase.

In addition to the above described $\Delta 6$ desaturation capability of mammalian FADS2, this enzyme can also operate as a $\Delta 8$ desaturase and thus PUFA substrates including 20:3n-3 and 20:2n-6 can be desaturated to 20:4n-3 and 20:3n-6, respectively, to thus reincorporate them into the pathway for further $\Delta 5$ desaturation [69] (Fig. 4). It is worth mentioning that the $\Delta 8$ desaturase capacity of the baboon FADS2 [69] seemed to be notably lower compared to those of marine teleost Fads2 assayed under similar conditions [70]. These findings represent further evidence of subfunctionalization occurring in teleost Fads2. Importantly, while the

teleostei Elovl5 has been identified to elongate C₁₈ PUFA (18:3n-3 and 18:2n-6) required for the $\Delta 8$ pathway, it still remains to be elucidated if the mammalian ELOVL5 is responsible for such conversions [22]

Adaptive evolution studies have recently revealed that ability to biosynthesize LC-PUFA varies notably among human populations and, for instance, levels of LC-PUFA in African-American individuals were found to be higher compared to European-Americans [158, 159]. Such differences were linked to polymorphisms in the FADS gene cluster that then lead to higher capacity to biosynthesis LC-PUFA from C₁₈ precursors. It was postulated that highly efficient desaturation alleles of *FADS* genes are particularly abundant in African populations where these were retained by positive selection driven by dietary limitation of preformed LC-PUFA in isolated regions [160].

3.4.1 Mammalian ELOVL

Seven members of the ELOVL protein family (ELOVL 1-7) have been described in mammals [20, 22]. The elongation of saturated and monounsaturated fatty acids in mammals can be catalyzed by four elongases, namely ELOVL1, ELOVL3, ELOVL6 and ELOVL7, that show different tissue distributions probably reflecting different functional roles. ELOVL1 is thought to be involved in the production of saturated fatty acids up to C₂₆ in length, as present in sphingolipids [161, 162]. It is expressed in most tissues, which may reflect a housekeeping role and a requirement for these fatty acids and lipids containing them in membranes, but it is also highly expressed in some parts of the central nervous system and may have an important role in the synthesis of sphingomyelin and hence myelin [162, 163]. ELOVL3 is suggested to control the synthesis of saturated and monounsaturated fatty acids of up to C₂₄ and is expressed mainly in brown and white adipose tissue, skin and liver [20, 164]. ELOVL3 is cold inducible in brown adipose tissue [164], displays a diurnal expression in the liver, and is regulated by gender-specific steroid hormones such as glucocorticoids, androgens and oestrogens [165]. ELOVL6 is involved in the elongation of C₁₂₋₁₆ saturated fatty acids up to C₁₈, but it has no capability to elongate beyond C₁₈. *ELOVL6* is ubiquitously expressed but especially in liver, brain, and other tissues with high lipid content such as white and brown adipose tissue [166, 167]. ELOVL7, the latest member of the ELOVL family to be identified, is highly expressed in the kidney, pancreas, adrenal glands and prostate [168]. Knockdown experiments of Elovl7 in carcinoma cell lines showed reduced levels of saturates 20:0, 22:0

and 24:0. Moreover, overexpression of Elov17 in microsomes has confirmed that Elov17 participates in elongation of saturated fatty acids with up to 24 carbons.

As with saturated and monounsaturated fatty acids, and as previously described for fish above, there are multiple enzymes for the elongation of PUFA in mammals that differ in tissue distributions and fatty acid specificity likely reflecting different functional roles. The *ELOVL2* and *ELOVL5* enzymes appear to be fairly ubiquitously expressed in most tissues in mammals although there can be species differences with *Elov15* in rats showing highest expression in lung and brain whereas human *ELOVL5* is particularly highly expressed in testis and adrenal gland that are characterized by relatively high levels of 22:5n-6 [141, 167] (Fig. 5). These two elongases have overlapping fatty acid substrate specificity with ELOVL5 being able to elongate both C₁₈ and C₂₀ PUFA, but with no activity towards C₂₂, whereas ELOVL2 is generally able to elongate C₂₀ and C₂₂ PUFA [142, 166]. However, while mouse ELOVL2 was able to elongate 18:3n-6 to some degree, human ELOVL2 was not, which suggested there may be some functional divergence between species as discussed above in teleosts (Section 3.2.2.2). Neither ELOVL2 nor ELOVL5 have activity towards saturated or monounsaturated fatty acids [142, 166], contrary to what was observed with some teleost Elov15 when expressed in yeast. The yeast expression system was also used to investigate the substrate specificities, competitive substrate interactions and dose response curves of the rat ELOVL2 and ELOVL5 [146]. It was confirmed that the rat ELOVL2 was active towards C₂₀ and C₂₂ PUFA but interestingly, the reaction converting the C₂₂ PUFA, 22:5n-3 to 24:5n-3 appeared to be saturated at substrate concentrations and thus ELOVL2 was postulated as an additional control point of LC-PUFA biosynthesis in addition to FADS2 mentioned above.

In contrast to ELOVL2 and ELOVL5, ELOVL4 has much more specific tissue distribution being highly expressed in retina and brain but also skin [169, 170]. A major difference of ELOVL4 compared to the other mammalian ELOVL is that it is capable of elongating both saturated fatty acids and PUFA [20]. Therefore, ELOVL4 is responsible for the biosynthesis of saturated VLC-FA (e.g. 26:0, 28:0 and 30:0) that are components of sphingolipids and ceramides and important in the skin [171, 172], and also for the biosynthesis of VLC-PUFA (> C₂₄) that are components of key phosphatidylcholine species in retina [173-175], brain [14, 176] and testis [177-180]. Although some of these studies revealed potential substrates for mammalian ELOVL4, the specific steps in which ELOVL4 was involved was only described in detail later when the human orthologue of ELOVL4 was characterized [181]. Overexpression of human ELOVL4 in rat neonatal cardiomyocytes and a human RPE cell line ARPE-19 showed a decreased level of 26:0 and concomitant increased levels of 28:0 and

30:0. Similar conclusions were obtained in genetically engineered mice lacking a functional ELOVL4 protein, which showed increased levels of C₂₆ fatty acids and a depletion of C₂₈ in lipids of the epidermal stratum corneum, which altered the skin barrier function and ultimately caused dehydration and perinatal death [171, 172, 182]. In addition to saturated VLC-FA biosynthesis, the human ELOVL4 was also demonstrated to mediate the production of C₂₈-C₃₈ n-3 VLC-PUFA, compounds mainly present into phospholipid molecules in retina [183]. As far as the authors are aware, no other mammalian ELOVL4 has been functionally characterized.

4. Concluding remarks

The extraordinary number of full genome sequences currently available offers a unique opportunity to investigate the evolution of complex gene pathways, and their impact on animal physiology. One of those examples is the genetic cascade controlling LC-PUFA biosynthesis, namely Fads and Elovl enzymes. In this review, we extensively addressed aspects directly linked with the diversity and function of Elovl and Fads genes in key chordate species. The existing literature supports the fundamental conclusion that the wiring and elaboration of a complete and functional LC-PUFA cascade evolved in the vertebrate lineage. The available data support an evolutionary model where a combination of whole genome and tandem duplications expanded the repertoire of Elovl and Fads in vertebrate ancestry, thus contributing to the diversification of these molecular modules – Elovl2, Elovl4, Elovl5, Fads1, and Fads2 (Fig. 7).

In this context, some questions remain to be addressed. The characterization of FADS enzymes in the amphioxus and agnathans will be essential to provide a defined picture at the invertebrate/vertebrate transition. Additionally, the functional characterization of Fads and Elovl genes retrieved from the sequenced genome of the holostean spotted gar, a basal freshwater fish closely related to teleosts, should facilitate our understanding of LC-PUFA biosynthesis. Also, Elovl4 orthologues remain largely uncharacterized as well their exact role in different species. Finally, the integration with functional *in vivo* studies to determine the interplay between endogenous synthesis versus dietary inputs of LC-PUFA is of vital importance to understand the impact and regulation at the physiological level.

Acknowledgements

LFCC research is funded by the Foundation for Science and Technology (Strategic Funding UID/Multi/04423/2013).

References

- [1] Schmitz G, Ecker J. The opposing effects of n-3 and n-6 fatty acids. *Prog Lipid Res* 2008; 47:147-55.
- [2] Delsuc F, Brinkmann H, Chourrout D, Philippe H. Tunicates and not cephalochordates are the closest living relatives of vertebrates. *Nature* 2006; 439:965-8.
- [3] Shimeld SM, Holland PW. Vertebrate innovations. *Proc Natl Acad Sci USA* 2000; 97:4449-52.
- [4] Braasch I, Scharl M. Evolution of endothelin receptors in vertebrates. *Gen Comp Endocrinol* 2014; 209:21-34.
- [5] Castro LF, Gonçalves O, Mazan S, Tay BH, Venkatesh B, Wilson JM. Recurrent gene loss correlates with the evolution of stomach phenotypes in gnathostome history. *Proc Biol Sci* 2014; 281:2669.
- [6] Putnam NH, Butts T, Ferrier DE, Furlong RF, Hellsten U, Kawashima T et al. The amphioxus genome and the evolution of the chordate karyotype. *Nature* 2008; 453:1064-71.
- [7] Kuraku S, Meyer A, Kuratani S. Timing of genome duplications relative to the origin of the vertebrates: did cyclostomes diverge before or after? *Mol Biol Evol* 2009; 26:47-59.
- [8] Mehta TK, Ravi V, Yamasaki S, Lee AP, Lian MM, Tay BH, et al. Evidence for at least six Hox clusters in the Japanese lamprey (*Lethenteron japonicum*). *Proc Natl Acad Sci USA* 2013; 110:16044-9.
- [9] Smith JJ, Keinath MC. The sea lamprey meiotic map improves resolution of ancient vertebrate genome duplications. *Genome Res* 2015; 25:1081-90.
- [10] Evans H, De Tomaso T, Quail M, Rogers J, Gracey AY, Cossins AR, Berenbrink M. Ancient and modern duplication events and the evolution of stearoyl-CoA desaturases in teleost fishes. *Physiol Genomics* 2008; 35:18-29.
- [11] Wakil SJ, Stoops JK, Joshi VC. Fatty acid synthesis and its regulation. *Ann Rev Biochem* 1983; 52:537-79.
- [12] Khosla C, Gokhale RS, Jacobsen JR, Cane DE. Tolerance and specificity of polyketide synthases. *Ann Rev Biochem* 1999; 68:219-53.

985 [13] Bell MV, Tocher DR. Biosynthesis of fatty acids; general principles and new directions.
986 In: Arts MT, Brett M, Kainz M, editors. *Lipids in Aquatic Ecosystems*, New York:
987 Springer-Verlag; 2009, p. 211-36.

988 [14] Poulos A. Very long chain fatty acids in higher animals – A review. *Lipids* 1995; 30:1-
989 14.

990 [15] Gunstone FD, Harwood JL. Occurrence and characteristics of oils and fats. In: Gunstone
991 FD, Harwood JL, Dijkstra AJ, editors. *The lipid handbook* (3rd edition), Boca Raton:
992 Taylor & Francis, CRC Press; 2007, p. 37-142.

993 [16] Lemieux B, Miguel M, Somerville C, Browse J. Mutants of *Arabidopsis* with alterations
994 in seed lipid fatty acid composition. *Theor Appl Genet* 1990; 80:234–40.

995 [17] Venegas-Calcrón M, Beaudoin F, Sayanova O, Napier JA. Co-transcribed genes for long
996 chain polyunsaturated fatty acid biosynthesis in the protozoon *Perkinsus marinus* include a
997 plant-like FAE1 3-Ketoacyl Coenzyme A Synthase. *J Biol Chem* 2007; 82:2996–3003.

998 [18] Harwood JL, Guschina IA. The versatility of algae and their lipid metabolism. *Biochimie*
999 2009; 91:679-84.

1000 [19] Leonard AE, Pereira SL, Sprecher H, Huang YS. Elongation of long-chain fatty acids.
1001 *Prog Lipid Res* 2004; 43:36-54.

1002 [20] Jakobsson A, Westerberg R, Jacobsson A. Fatty acid elongases in mammals: Their
1003 regulation and roles in metabolism. *Prog Lipid Res* 2006; 45:237-49.

1004 [21] Parker-Barnes JM, Das T, Bobik E, Leonard AE, Thurmond JM, Chuang L-T, Huang YS,
1005 Mukerji P. Identification and characterization of an enzyme involved in the elongation of
1006 n-6 and n-3 polyunsaturated fatty acids. *Proc Natl Acad Sci USA* 2000; 97:8284–8289.

1007 [22] Guillou H, Zdravce D, Martin PGP, Jacobsson A. The key roles of elongases and
1008 desaturases in mammalian fatty acid metabolism: Insights from transgenic mice. *Prog*
1009 *Lipid Res* 2010; 49:186-99.

1010 [23] Moon YA, Horton JD. Identification of two mammalian reductases involved in the two-
1011 carbon fatty acyl elongation cascade. *J Biol Chem* 2003; 278:7335–43.

1012 [24] Naganuma T, Sato Y, Sassa T, Ohno Y, Kihara A. Biochemical characterization of the
1013 very long-chain fatty acid elongase ELOVL7. *FEBS Lett* 2011; 585:3337–41.

1014 [25] Venegas-Calcrón M, Sayanova O, Napier JA. An alternative to fish oils: Metabolic
1015 engineering of oil-seed crops to produce omega-3 long chain polyunsaturated fatty acids.
1016 *Prog Lipid Res* 2010; 49:108–119.

1017 [26] Haslam TM, Kunst L. Extending the story of very-long-chain fatty acid elongation. *Plant*
1018 *Sci* 2013; 210: 93-107.

1019 [27] Orikasa Y, Yamada A, Yu R, Ito Y, Nishida T, Yumoto I, et al. Characterization of the
1020 eicosapentaenoic acid biosynthesis gene cluster from *Shewanella* sp strain SCRC-2738.
1021 Cell Mol Biol 2004; 50:625-30.

1022 [28] Allen EE, Bartlett DH. Structure and regulation of the omega-3 polyunsaturated fatty
1023 acid synthase genes from the deep-sea bacterium *Photobacterium profundum* strain SS9.
1024 Microbiol 2002; 148:1903–13.

1025 [29] Tanaka M, Ueno A, Kawasaki K, Yumoto I, Ohgiya S, Hoshino T, et al. Isolation of
1026 clustered genes that are notably homologous to the eicosapentaenoic acid biosynthesis
1027 gene cluster from the docosahexaenoic acid-producing bacterium *Vibrio marinus* strain
1028 MP-1. Biotechnol Lett 1999; 21:939-45.

1029 [30] Metz JG, Roessler P, Facciotti D, Levering C, Dittrich F, Lassner M, et al. Production of
1030 polyunsaturated fatty acids by polyketide synthases in both prokaryotes and eukaryotes.
1031 Science 2001; 293:290-3.

1032 [31] Nakamura MT, Nara TY. Structure, function, and dietary regulation of delta6, delta5,
1033 and delta9 desaturases. Annu Rev Nutr 2004; 24:345–76.

1034 [32] Meesapyodsuk D, Qiu X. The front-end desaturase: Structure, function, evolution and
1035 biotechnological use. Lipids 2012; 47:227-237.

1036 [33] Shanklin J, Cahoon EB. Desaturation and related modifications of fatty acids. Annu Rev
1037 Plant Physiol Plant Mol Biol 1998; 49:611–641.

1038 [34] Sperling P, Ternes P, Zank TK, Heinz E. The evolution of desaturases. Prostaglandins
1039 Leukot Essent Fatty Acids 2003; 68:73–95.

1040 [35] Hashimoto K, Yoshizawa AC, Saito K, Yamada T, Kanehisa M. The repertoire of
1041 desaturases for unsaturated fatty acid synthesis in 397 genomes. Genome Inform 2006;
1042 17:173–83.

1043 [36] Zhu K, Choi KH, Schweizer HP, Rock CO, Zhang YM. Two aerobic pathways for the
1044 formation of unsaturated fatty acids in *Pseudomonas aeruginosa*. Mol Microbiol 2006;
1045 60:260–73.

1046 [37] Yasunaga K, Shiro Y, Yoshinori N. The occurrence of direct desaturation of
1047 phospholipid acyl chain in *Tetrahymena pyriformis*: Thermal adaptation of membrane
1048 phospholipid. Biochim Biophys Acta 1980; 618:214-22.

1049 [38] Tocher DR, Leaver MJ, Hodgson PA. Recent advances in the biochemistry and
1050 molecular biology of fatty acyl desaturases. Prog Lipid Res 1998; 37:73-117.

- [39] Lopez Alonso D, Garcia-Maroto F, Rodriguez-Ruiz J, Garrido JA, Vilches MA. Evolution of the membrane-bound fatty acid desaturases. *Biochem Systematics Ecol* 2003; 31:1111–24.
- [40] Gostincar C, Turk M, Gunde-Cimerman N. The evolution of fatty acid desaturases and cytochrome b₅ in eukaryotes. *J Membrane Biol* 2010; 233, 63–72.
- [41] Napier JA, Hey SJ, Lacey DJ, Shewry PR. Identification of a *Caenorhabditis elegans* $\Delta 6$ fatty acid - desaturase by heterologous expression in *Saccharomyces cerevisiae*. *Biochem J* 1998; 330:611-14.
- [42] Harwood JL. Recent advances in the biosynthesis of plant fatty acids. *Biochim Biophys Acta* 1996; 1301:7-56.
- [43] Martin CE, Oh CS, Kandasamy P, Chellapa R, Vemula M. Yeast desaturases. *Biochem Soc. Trans.* 2002; 30:1080–2.
- [44] Dobrzyn A, Ntambi JM. The role of stearoyl-CoA desaturase in the control of metabolism. *Prostaglandins Leukot Essent Fatty Acids* 2005; 73:35–41.
- [45] Bai Y, McCoy JG, Levin EJ, Sobrado P, Rajashankar KR, Fox BG, Zhou M. X-ray structure of a mammalian stearoyl-CoA desaturase. *Nature* 2015; 524: 252–6.
- [46] Wang H, Klein MG, Zou H, Lane W, Snell G, Levin I, et al. Crystal structure of human stearoyl-coenzyme A desaturase in complex with substrate. *Nature Struct Mol Biol* 2015; 22:581-5.
- [47] Castro LFC, Wilson JM, Gonçalves O, Galante-Oliveira S, Rocha E, Cunha I. The evolutionary history of the stearoyl-CoA desaturase gene family in vertebrates. *BMC Evol Biol* 2011; 11:132.
- [48] Guschina IA, Harwood JL. Lipids and lipid metabolism in eukaryotic algae. *Prog Lipid Res* 2006; 45:160-86.
- [49] Chi X, Yang Q, Zhao F, Qin S, Yang Y, Shen J, Lin H. Comparative analysis of fatty acid desaturases in Cyanobacterial genomes. *Comp Funct Genomics* 2008; 284508.
- [50] Kotajima T, Shiraiwa Y, Suzuki I. Functional screening of a novel Delta15 fatty acid desaturase from the coccolithophorid *Emiliania huxleyi*. *Biochim Biophys Acta* 2014; 1842:1451-8.
- [51] Pereira SL, Leonard AE, Mukerji P. Recent advances in the study of fatty acid desaturases from animals and lower eukaryotes. *Prostaglandins Leukot Essent Fatty Acids* 2003; 68:97-106.
- [52] Jones AL, Lloyd D, Harwood JL. Rapid induction of microsomal delta 12(omega 6)-desaturase activity in chilled *Acanthamoeba castellanii*. *Biochem J* 1993; 296:183-8.

1085 [53] Sayanova O, Haslam R, Guschina I, Lloyd D, Christie WW, Harwood, JL, Napier JA. A
1086 bifunctional $\Delta 12, \Delta 15$ -desaturase from *Acanthamoeba castellanii* directs the synthesis of
1087 highly unusual n-1 series unsaturated fatty acids. J Biol Chem 2006; 281:36533–41.

1088 [54] Zhang S, Sakuradani E, Ito K, Shimizu S. Identification of a novel bifunctional $\Delta 12 / \Delta 15$
1089 fatty acid desaturase from a basidiomycete, *Coprinus cinereus* TD#822-2. FEBS Lett
1090 2007; 581:315-9.

1091 [55] Borgeson CE, de Renobales M Blomquist GJ. Characterization of the $\Delta 12$ desaturase in
1092 the American cockroach, *Periplaneta americana*: the nature of the substrate. Biochim
1093 Biophys Acta 1990; 1047:135–40.

1094 [56] Cripps C, Borgeson C, Blomquist GJ, de Renobales M. The $\Delta 12$ -desaturase from the
1095 house cricket *Acheta domesticus* (Orthoptera: Gryllidae): characterization and form of the
1096 substrate. Arch Biochem Biophys 1990; 278:46–51.

1097 [57] Zhou XR, Horne I, Damcevski K, Haritos V, Green A, Singh S. Isolation and functional
1098 characterization of two independently-evolved fatty acid $\Delta 12$ -desaturase genes from
1099 insects. Insect Mol Biol 2008; 17:667–76.

1100 [58] Vanhercke T, Shresth P, Green AG, Singh SP. Mechanistic and structural insights into
1101 the regioselectivity of an acyl-CoA fatty acid desaturase via directed molecular evolution. J
1102 Biol Chem 2011; 286:12860-9.

1103 [59] Spychalla JP, Kinney AJ, Browse J. Identification of an animal omega-3 fatty acid
1104 desaturase by heterologous expression in Arabidopsis. Proc Natl Acad Sci USA 1997;
1105 94:1142–7.

1106 [60] Peyou-Ndi MM, Watts JL, Browse J. Identification and characterization of an animal Δ
1107 (12) fatty acid desaturase gene by heterologous expression in *Saccharomyces cerevisiae*.
1108 Arch Biochem Biophys 2000; 376:399–408.

1109 [61] Zhou XR, Green AG, Singh SP. *Caenorhabditis elegans* $\Delta 12$ -desaturase FAT-2 is a
1110 bifunctional desaturase able to desaturate a diverse range of fatty acid substrates at the $\Delta 12$
1111 and $\Delta 15$ positions. J Biol Chem 2011; 286:43644–50.

1112 [62] Aitzetmuller K, Tsevegsuren N. Seed fatty acid “front-end” desaturases and
1113 chemotaxonomy - a case study in the Ranunculaceae. J Plant Physiol 1994; 143:538–43.

1114 [63] Okuyama H, Orikasa Y, Nishida T, Watanabe K, Morita N. Bacterial genes responsible
1115 for the biosynthesis of eicosapentaenoic and docosahexaenoic acids and their heterologous
1116 expression. Appl Environmental Microbiol 2007; 73: 665–70.

1117 [64] Knutzon DS, Thurmond JM, Huang Y-S, Chaudhary S, Bodik EG, Chan GM, et al.
1118 Identification of $\Delta 5$ -desaturase from *Mortierella alpina* by heterologous expression in
1119 bakers' yeast and canola. J Biol Chem 1998; 273:29360-6.

1120 [65] Sakuradani E, Kobayashi M, Shimizu S. Fatty acid desaturase from an arachidonic acid-
1121 producing *Mortierella* fungus: Gene cloning and its heterologous expression in a fungus,
1122 *Aspergillus*. Gene 1999; 238:445-53.

1123 [66] Sakuradani E, Shimizu S. Gene cloning and functional analysis of a second $\Delta 6$ -Fatty acid
1124 desaturase from an arachidonic acid- producing *Mortierella* fungus. Biosci Biotechnol
1125 Biochem 2003; 67:704-11.

1126 [67] Lu H, Li J-N, Chai Y-R, Zhang X-K. Identification and characterization of a novel $\Delta 6$ -
1127 fatty acid desaturase gene from *Rhizopus nigricans*. Mol Biol Rep 2009; 36:2291-7.

1128 [68] Cook HW, McMaster CR. Fatty acid desaturation and chain elongation in eukaryotes. In:
1129 Vance DE, Vance JE, editors. Biochemistry of Lipids, Lipoproteins and Membranes (4th
1130 Edition), Amsterdam: Elsevier; 2002, p. 181-204.

1131 [69] Park WJ, Kothapalli KS, Lawrence P, Tybureczy C, Brenna JT. An alternate pathway to
1132 long-chain polyunsaturates: The FADS2 gene product $\Delta 8$ -desaturates 20:2n-6 and
1133 20:3n-3. J Lipid Res 2009; 50:1195-202.

1134 [70] Monroig Ó, Li Y, Tocher DR. $\Delta 8$ desaturation activity varies among fatty acyl
1135 desaturases of teleost fish: high activity in $\Delta 6$ desaturases of marine species. Comp
1136 Biochem Physiol Part B 2011; 159:206-213.

1137 [71] Li Y, Monroig Ó, Zhang L, Wang S, Zheng X, Dick JR, et al. Vertebrate fatty acyl
1138 desaturase with $\Delta 4$ activity. Proc Natl Acad Sci USA 2010; 107:16840-5.

1139 [72] Sprecher H. Metabolism of highly unsaturated n-3 and n-6 fatty acids. Biochim Biophys
1140 Acta 2000; 1486:219-31.

1141 [73] Park HG, Park WJ, Kothapalli KS, Brenna JT. The fatty acid desaturase 2 (FADS2) gene
1142 product catalyzes $\Delta 4$ desaturation to yield n-3 docosahexaenoic acid and n-6
1143 docosapentaenoic acid in human cells. FASEB J 2015; 29:3911-9.

1144 [74] Hui JHL, McDougall C, Monteiro AS, Holland PWH, Arendt D, Balavoine G, Ferrier
1145 DEK. Extensive chordate and annelid macrosynteny reveals ancestral homeobox gene
1146 organization. Mol Biol Evol 2012; 29:157-65.

1147 [75] Raible F, Tessmar-Raible K, Osoegawa K, Wincker P, Jubin C, Balavoine G et al.
1148 Vertebrate-type intron-rich genes in the marine annelid *Platynereis dumerilii*. Science
1149 2005; 310: 1325-26.

1150 [76] Takahashi T, McDougall C, Troscianko J, Chen W-C, Jayaraman-Nagarajan A, Shimeld
1151 SM, Ferrier DEK. An EST screen from the annelid *Pomatoceros lamarckii* reveals patterns
1152 of gene loss and gain in animals. BMC Evol Biol 2009;9:240.

1153 [77] Monroig Ó, Tocher DR, Navarro JC. Biosynthesis of polyunsaturated fatty acids in
1154 marine invertebrates: Recent advances in molecular mechanisms. Mar Drugs 2013;
1155 11:3998-4018.

1156 [78] Michaelson LV, Napier JA, Lewis M, Griffiths G, Lazarus CM, Stobart AK. Functional
1157 identification of a fatty acid $\Delta 5$ desaturase gene from *Caenorhabditis elegans*. FEBS Lett
1158 1998. 439:215-218.

1159 [79] Watts JL, Browse J. Isolation and characterisation of a $\Delta 5$ fatty acid desaturase from
1160 *Caenorhabditis elegans*. Arch Biochem Biophys 1999; 362:175-82.

1161 [80] Monroig Ó, Navarro JC, Dick JR, Alemany F, Tocher DR. Identification of a $\Delta 5$ -like
1162 fatty acyl desaturase from the cephalopod *Octopus vulgaris* (Cuvier 1797) involved in the
1163 biosynthesis of essential fatty acids. Mar Biotech 2012; 14:411–22.

1164 [81] Monroig Ó, Guinot D, Hontoria F, Tocher DR, Navarro JC. Biosynthesis of essential
1165 fatty acids in *Octopus vulgaris* (Cuvier, 1797): Molecular cloning, functional
1166 characterisation and tissue distribution of a fatty acyl elongase. Aquaculture 2012; 360-
1167 361:45-53.

1168 [82] Li M, Mai K, He G, Ai Q, Zhang W, Xu W, et al. Characterization of $\Delta 5$ fatty acyl
1169 desaturase in abalone *Haliotis discus hannai* Ino. Aquaculture 2013; 416–417:48–56.

1170 [83] Liu H, Zheng H, Wang S, Wang Y, Li S, Liu W, Zhang G. Cloning and functional
1171 characterization of a polyunsaturated fatty acid elongase in a marine bivalve noble scallop
1172 *Chlamys nobilis* Reeve. Aquaculture 2013; 416: 146–51.

1173 [84] Liu H, Guo Z, Zheng H, Wang S, Wang Y, Liu W, Zhang G. Functional characterization
1174 of a $\Delta 5$ -like fatty acyl desaturase and its expression during early embryogenesis in the
1175 noble scallop *Chlamys nobilis* Reeve. Mol Biol Rep 2014; 41:7437–45.

1176 [85] Monroig Ó, Hontoria F, Varó I, Tocher DR, Navarro JC. Investigating the essential fatty
1177 acids in the common cuttlefish *Sepia officinalis* (Mollusca, Cephalopoda): Molecular
1178 cloning and functional characterisation of fatty acyl desaturase and elongase. Aquaculture
1179 2016; 450:38–47.

1180 [86] Surm JM, Prentis PJ, Pavasovic A. Comparative analysis and distribution of Omega-3
1181 LC-PUFA biosynthesis genes in marine molluscs. PLoS ONE 2015; 10:e0136301.

- [87] Monroig O, Marquez-Lopes M, Navarro JC, Hontoria F, Ruivo R, Santos MM, et al. Evolutionary functional elaboration of the Elovl2/5 gene family in chordates. *Sci Rep* (in press).
- [88] Meyer A, Kirsch H, Domergue F, Abbadi A, Sperling P, Bauer J, et al. Novel fatty acid elongases and their use for the reconstitution of docosahexaenoic acid biosynthesis. *J Lipid Res* 2004; 45:1899-909.
- [89] Castro LFC, Monroig Ó, Leaver MJ, Wilson J, Cunha I, Tocher DR. Functional desaturase Fads1 ($\Delta 5$) and Fads2 ($\Delta 6$) orthologues evolved before the origin of jawed vertebrates. *PLoS ONE* 2012; 7:e31950.
- [90] Gregory M, Geier MS, Gibson RA, James MJ. Functional characterization of the chicken fatty acid elongases. *J Nutr* 2013; 143:12-6.
- [91] Gregory M, James MJ. Functional characterization of the duck and turkey fatty acyl elongase enzymes ELOVL5 and ELOVL2. *J Nutr* 2014; 11:1234-9.
- [92] Fonseca-Madrigal J, Navarro JC, Hontoria F, Tocher DR, Martínez-Palacios CA, Monroig Ó. Diversification of substrate specificities in teleostei Fads2: Characterization of $\Delta 4$ and $\Delta 6\Delta 5$ desaturases of *Chirostoma estor*. *J Lipid Res* 2014;55:1408-19.
- [93] Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, et al. Elephant shark genome provides unique insights into gnathostome evolution. *Nature* 2014; 505:174-9.
- [94] Noro M, Sugahara F, Kuraku S. Reevaluating Emx gene phylogeny: homopolymeric amino acid tracts as a potential factor obscuring orthology signals in cyclostome genes. *BMC Evol Biol* 2015; 15:78.
- [95] Tocher DR. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquacult Res* 2010; 41:717–32.
- [96] Carmona-Antoñanzas G, Tocher DR, Taggart JB, Leaver MJ. An evolutionary perspective on Elovl5 fatty acid elongase: comparison of Northern pike and duplicated paralogs from Atlantic salmon. *BMC Evol Biol* 2013; 13:85.
- [97] Morais S, Castanheira F, Martínez-Rubio L, Conceição LEC, Tocher DR. Long-chain polyunsaturated fatty acid synthesis in a marine vertebrate: ontogenetic and nutritional regulation of a fatty acyl desaturase with $\Delta 4$ activity. *Biochim Biophys Acta* 2012; 1821:660–71.
- [98] Monroig Ó, Tocher DR, Hontoria F, Navarro JC. Functional characterisation of a Fads2 fatty acyl desaturase with $\Delta 6/\Delta 8$ activity and an Elovl5 with C16, C18 and C20 elongase activity in the anadromous teleost meagre (*Argyrosomus regius*). *Aquaculture* 2013; 412-413:14–22.

1216 [99] Seiliez I, Panserat S, Kaushik S, Bergot P. Cloning, tissue distribution and nutritional
1217 regulation of a $\Delta 6$ -desaturase-like enzyme in rainbow trout. *Comp Biochem Physiol Part B*
1218 2001; 130:83–93.

1219 [100] Hastings N, Agaba M, Tocher DR, Leaver MJ, Dick JR, Sargent JR, Teale AJ. A
1220 vertebrate fatty acid desaturase with $\Delta 5$ and $\Delta 6$ activities. *Proc Natl Acad Sci USA* 2001;
1221 98:14304-9.

1222 [101] Zheng X, Seiliez I, Hastings N, Tocher DR, Panserat S, Dickson CA, et al.
1223 Characterization and comparison of fatty acyl $\Delta 6$ desaturase cDNAs from freshwater and
1224 marine teleost fish specie. *Comp Biochem Physiol Part B* 2004; 139:269-79.

1225 [102] Hastings N, Agaba MK, Tocher DR, Zheng X, Dickson CA, Dick JR, Teale AJ.
1226 Molecular cloning and functional characterization of fatty acyl desaturase and elongase
1227 cDNAs involved in the production of eicosapentaenoic and docosahexaenoic acids from α -
1228 linolenic acid in Atlantic salmon (*Salmo salar*). *Mar Biotechnol* 2005; 6:463-74.

1229 [103] Zheng X, Tocher DR, Dickson CA, Dick JR, Bell JG, Teale AJ. Highly unsaturated
1230 fatty acid synthesis in vertebrates: new insights with the cloning and characterisation of a
1231 $\Delta 6$ desaturase of Atlantic salmon. *Lipids* 2005; 40:13-24.

1232 [104] Monroig Ó, Zheng X, Morais S, Leaver MJ, Taggart JB, Tocher DR. Multiple genes for
1233 functional $\Delta 6$ fatty acyl desaturases (Fad) in Atlantic salmon (*Salmo salar* L.): Gene and
1234 cDNA characterization, functional expression, tissue distribution and nutritional regulation.
1235 *Biochim Biophys Acta* 2010; 1801:1072–81.

1236 [105] Seiliez I, Panserat S, Corraze G, Kaushik S, Bergot P. Cloning and nutritional
1237 regulation of a $\Delta 6$ -desaturase-like enzyme in the marine teleost gilthead sea bream (*Sparus*
1238 *aurata*). *Comp Biochem Physiol Part B* 2003; 135:449–60.

1239 [106] Tocher DR, Zheng X, Schlechtriem C, Hastings N, Dick JR, Teale AJ. Highly
1240 unsaturated fatty acid synthesis in marine fish: cloning, functional characterization and
1241 nutritional regulation of fatty acid $\Delta 6$ desaturase of Atlantic cod (*Gadus morhua* L.).
1242 *Lipids* 2006; 41:1003-16.

1243 [107] Zheng X, Ding Z, Xu Y, Monroig O, Morais S, Tocher DR. Physiological roles of fatty
1244 acyl desaturase and elongase in marine fish: Characterisation of cDNAs of fatty acyl $\Delta 6$
1245 desaturase and Elovl5 elongase of cobia (*Rachycentron canadum*). *Aquaculture* 2009;
1246 290:122-31.

1247 [108] González-Rovira A, Mourente G, Zheng X, Tocher DR, Pendón C. Molecular and
1248 functional characterization and expression analysis of a $\Delta 6$ fatty acyl desaturase cDNA of
1249 European sea bass (*Dicentrarchus labrax* L.). *Aquaculture* 2009; 298:90–100.

1250 [109] Santigosa E, Geay F, Tonon T, Le Delliou H, Kuhl H, Reinhardt R et al. Cloning, tissue
 1251 expression analysis, and functional characterization of two $\Delta 6$ -desaturase variants of sea
 1252 bass (*Dicentrarchus labrax* L.). Mar Biothechnol 2011; 13:22-31.
 1253 [110] Mohd-Yusof NY, Monroig Ó, Mohd-Adnan A, Wan K-L, Tocher DR. Investigation of
 1254 highly unsaturated fatty acid metabolism in the Asian sea bass, *Lates calcarifer*. Fish
 1255 Physiol Biochem 2010; 36:827-43.
 1256 [111] Kim SH, Kim JB, Kim SY, Roh KH, Kim HU, Lee K-R, et al. Functional
 1257 characterization of a delta 6-desaturase gene from the black seabream (*Acanthopagrus*
 1258 *schlegeli*). Biotechnol Lett 2011; 33:1185–93.
 1259 [112] Kabeya N, Yamamoto Y, Cummins SF, Elizur A, Yazawa R, Takeuchi Y, Haga Y,
 1260 Satoh S, Yoshizaki G. Polyunsaturated fatty acid metabolism in a marine teleost, Nibe
 1261 croaker *Nibea mitsukurii*: Functional characterization of Fads2 desaturase and Elovl5 and
 1262 Elovl4 elongases. Comp Biochem Physiol Part B 2015; 188:37–45.
 1263 [113] Morais S, Mourente G, Ortega A, Tocher JA, Tocher DR. Expression of fatty acyl
 1264 desaturase and elongase genes, and evolution of DHA:EPA ratio during development of
 1265 Atlantic bluefin tuna (*Thunnus thynnus* L.). Aquaculture 2011; 313:129-39.
 1266 [114] Tanomman S, Ketudat-Cairns M, Jangprai A, Boonanuntanasarn S. Characterization of
 1267 fatty acid delta-6 desaturase gene in Nile tilapia and heterogenous expression in
 1268 *Saccharomyces cerevisiae*. Comp Biochem Physiol Part B 2013; 166:148-56.
 1269 [115] Wang S, Monroig Ó, Zhang L, You C, Tocher DR, Li Y. Investigating the long-chain
 1270 polyunsaturated fatty acid biosynthesis in teleost fish: Functional characterisation of fatty
 1271 acyl desaturase (Fads2) and Elovl5 elongase in the catadromous species, Japanese eel
 1272 *Anguilla japonica*. Aquaculture 2014; 434:57–65.
 1273 [116] Characterization, mRNA expression and regulation of $\Delta 6$ fatty acyl desaturase
 1274 (FADS2) by dietary n – 3 long chain polyunsaturated fatty acid (LC-PUFA) levels in
 1275 grouper larvae (*Epinephelus coioides*). Aquaculture 2014; 434:212–9.
 1276 [117] Kuah M-K, Jaya-Ram A, Shu-Chien AC. The capacity for long-chain polyunsaturated
 1277 fatty acid synthesis in a carnivorous vertebrate: Functional characterisation and nutritional
 1278 regulation of a Fads2 fatty acyl desaturase with $\Delta 4$ activity and an Elovl5 elongase in
 1279 striped snakehead (*Channa striata*). Biochim Biophys Acta 2015; 1851:248–60.
 1280 [118] Buzzi M, Henderson RJ, Sargent JR. The desaturation and elongation of linolenic acid
 1281 and eicosapentaenoic acid by hepatocytes and liver microsomes from rainbow trout
 1282 (*Oncorhynchus mykiss*) fed diets containing fish oil or olive oil. Biochim Biophys Acta
 1283 1996; 1299:235-44.

- [119] Buzzi M, Henderson RJ, Sargent JR. Biosynthesis of docosahexaenoic acid in trout hepatocytes proceeds via 24-carbon intermediates. *Comp Biochem Physiol Part B* 1997; 116:263–7.
- [120] Tocher DR, Agaba M, Hastings N, Teale AJ. Biochemical and molecular studies of the fatty acid desaturation pathway in fish. In: Browman HI, Skiftesvik AB, editors. *The Big Fish Bang – Proceedings of the 26th Annual Larval Fish Conference*, Bergen: Institute of Marine Nutrition; 2003, p. 211-27.
- [121] Betancur-R R, Broughton RE, Wiley EO, Carpenter K, López JA, Li C, et al. The tree of life and a new classification of bony fishes. *PLoS Curr.* 2013; 18:1-45. doi: 10.1371/currents.tol.53ba26640df0ccae75bb165c8c26288
- [122] Morais S, Mourente G, Martínez A, Gras N, Tocher DR. Docosahexaenoic acid biosynthesis via fatty acyl elongase and $\Delta 4$ -desaturase and its modulation by dietary lipid level and fatty acid composition in a marine vertebrate. *Biochim Biophys Acta* 2015; 1851:588–97.
- [123] Ferdinandusse S, Denis S, van Roermund CWT, Wanders RJA, Dacremont G. Identification of the peroxisomal-oxidation enzymes involved in the degradation of long-chain dicarboxylic acids. *J Lipid Res* 2004; 45:1104–11.
- [124] Lim ZL, Senger T, Vrinten P. Four amino acid residues influence the substrate chain-length and regioselectivity of *Siganus canaliculatus* D4 and D5/6 desaturases. *Lipids* 2014; 49:357–67.
- [125] Damude HG, Zhang H, Farrall L, Ripp KG, Tomb J-F, Hollerbach D, Yadav NS. Identification of bifunctional $\Delta 12/\omega 3$ fatty acid desaturases for improving the ratio of $\omega 3$ to $\omega 6$ fatty acids in microbes and plants. *Proc Natl Acad Sci USA* 2006; 103:9446–51.
- [126] Serra M, Gauthier LT, Fabrias G, Buist PH. $\Delta 11$ desaturases of *Trichoplusia ni* and *Spodoptera littoralis* exhibit dual catalytic behaviour. *Insect Biochem Mol Biol* 2006; 36:822–25.
- [127] De Antueno RJ, Knickle LC, Smith H, Elliot ML, Allen SJ, Nwaka S, Winther MD. Activity of human $\Delta 5$ and $\Delta 6$ desaturases on multiple n-3 and n-6 polyunsaturated fatty acids. *FEBS Lett* 2001; 509:77-80.
- [128] D’Andrea S, Guillou H, Jan S, Catheline D, Thibault JN, Bouriel M, et al. The same rat $\Delta 6$ -desaturase not only acts on 18- but also on 24-carbon fatty acids in very-long-chain polyunsaturated fatty acid biosynthesis. *Biochem J* 2002; 364:49–55.

1316 [129] Leaver MJ, Bautista JM, Björnsson T, Jönsson E, Krey G, Tocher DR, Torstensen BE.
1317 Towards fish lipid nutrigenomics: current state and prospects for fin-fish aquaculture. *Rev*
1318 *Fisheries Sci* 2008; 16:71-92.

1319 [130] Agaba M, Tocher DR, Dickson C, Dick JR, Teale AJ. Zebrafish cDNA encoding
1320 multifunctional fatty acid elongase involved in production of eicosapentaenoic (20:5n-3)
1321 and docosahexaenoic (22:6n-3) acids. *Mar Biotechnol* 2004; 6:251-61.

1322 [131] Monroig Ó, Rotllant J, Sánchez E, Cerdá-Reverter JM, Tocher DR. Expression of long-
1323 chain polyunsaturated fatty acid (LC-PUFA) biosynthesis genes during zebrafish *Danio*
1324 *rerio* early embryogenesis. *Biochim Biophys Acta* 2009; 1791:1093–101.

1325 [132] Monroig Ó, Rotllant J, Cerdá-Reverter JM, Dick JR, Figueras A, Tocher DR.
1326 Expression and role of Elovl4 elongases in biosynthesis of very long-chain fatty acids
1327 during zebrafish *Danio rerio* early embryonic development. *Biochim Biophys Acta* 2010;
1328 1801:1145-54.

1329 [133] Agaba MK, Tocher DR, Dickson CA, Zheng X, Dick JR, Teale AJ. Cloning and
1330 functional characterisation of polyunsaturated fatty acid elongases from marine and
1331 freshwater teleost fish. *Comp Biochem Physiol Part B* 2005; 142:342–52.

1332 [134] Morais S, Monroig Ó, Zheng X, Leaver MJ, Tocher DR. Highly unsaturated fatty acid
1333 synthesis in Atlantic salmon: characterization of Elovl5- and Elovl2-like elongases. *Mar*
1334 *Biotechnol* 2009; 11:627–39.

1335 [135] Carmona-Antoñanzas G, Monroig Ó, Dick JR, Davie A, Tocher DR. Biosynthesis of
1336 very long-chain fatty acids (C>24) in Atlantic salmon: Cloning, functional characterisation,
1337 and tissue distribution of an Elovl4 elongase. *Comp Biochem Physiol Part B* 2011;
1338 159:122-9.

1339 [136] Monroig Ó, Webb K, Ibarra-Castro L, Holt GJ, Tocher DR. Biosynthesis of long-chain
1340 polyunsaturated fatty acids in marine fish: Characterization of an Elovl4-like elongase
1341 from cobia *Rachycentron canadum* and activation of the pathway during early life stages.
1342 *Aquaculture* 2011; 312:145–53.

1343 [137] Gregory M, See VHL, Gibson RA, Shuller KA. Cloning and functional characterisation
1344 of a fatty acyl elongase from southern bluefin tuna (*Thunnus maccoyii*). *Comp Biochem*
1345 *Physiol Part B* 2010; 155:178-85.

1346 [138] Monroig Ó, Wang S, Zhang L, You C, Tocher DR, Li Y. Elongation of long-chain fatty
1347 acids in rabbitfish *Siganus canaliculatus*: Cloning, functional characterisation and tissue
1348 distribution of Elovl5- and Elovl4-like elongases. *Aquaculture* 2012 ;350–353:63–70.

- [139] Kim SH, Kim JB, Jang YS, Kim SY, Roh KH, Kim HU, et al. Isolation and functional characterization of polyunsaturated fatty acid elongase (AsELOVL5) gene from black seabream (*Acanthopagrus schlegelii*). *Biotechnol Lett* 2012; 34:261–8.
- [140] Gregory MK, James MJ. Rainbow trout (*Oncorhynchus mykiss*) Elovl5 and Elovl2 differ in selectivity for elongation of omega-3 docosapentaenoic acid. *Biochim Biophys Acta* 2014; 1841: 1656–60.
- [141] Leonard AE, Bobik EG, Dorado J, Kroeger PE, Chuang L-T, Thurmond JM, et al. Cloning of a human cDNA encoding a novel enzyme involved in the elongation of long chain polyunsaturated fatty acids. *Biochem J* 2000; 350:765-70.
- [142] Leonard AE, Kelder B, Bobik EG, Chuang L-T, Lewis CJ, Kopchick JJ, et al. Identification and expression of mammalian long-chain PUFA elongation enzymes. *Lipids* 2002; 37:733-40.
- [143] Buzzi M, Henderson RJ, Sargent JR. The biosynthesis of docosahexaenoic acid [22:6(n-3)] from linolenic acid in primary hepatocytes isolated from wild northern pike. *J Fish Biol* 1997; 51: 1197-1208.
- [144] Carmona-Antoñanzas G, Zheng X, Tocher DR, Leaver MJ. Transposon insertions drive regulatory divergence of duplicated salmonid genes: Characterisation of *elovl5* homeologues with a major role in highly unsaturated fatty acid biosynthesis. *Mobile DNA* (under review).
- [145] Xue X, Feng CY, Hixson SM, Johnstone K, Anderson DM, Parrish CC, Rise ML. Characterization of the fatty acyl elongase (*elovl*) gene family, and hepatic *elovl* and delta-6 fatty acyl desaturase transcript expression and fatty acid responses to diets containing camelina oil in Atlantic cod (*Gadus morhua*). *Comp Biochem Physiol Part B* 2014; 175:9–22.
- [146] Gregory MK, Gibson RA, Cook-Johnson RJ, Cleland LG, James MJ. Elongase reactions as control points in long-chain polyunsaturated fatty acid synthesis. *PLoS ONE* 2011; 6: e29662.
- [147] Marquardt A, Stöhr H, White K, Weber BH. cDNA cloning, genomic structure, and chromosomal localization of three members of the human fatty acid desaturase family. *Genomics* 2000; 66:175-83.
- [148] Park WJ, Kothapalli KS, Reardon HT, Kim LY, Brenna JT. Novel fatty acid desaturase 3 (*FADS3*) transcripts generated by alternative splicing. *Gene* 2009; 446:28–34.
- [149] Blanchard H, Legrand P, Pédrone F. Fatty Acid Desaturase 3 (*Fads3*) is a singular member of the *Fads* cluster. *Biochimie* 2011; 93:87-90.

1383 [150] Reddy AS, Nuccio ML, Gross LM, Thomas TL. Isolation of a delta 6-desaturase gene
1384 from the *Cyanobacterium synechocystis* sp. strain PCC 6803 by gain-of-function
1385 expression in *Anabaena* sp. strain PCC 7120. Plant Mol Biol 1993; 22:293–300.

1386 [151] Sayanova O, Smith MA, Lapinskas P, Stobart AK, Dobson G, Christie WW, et al.
1387 Expression of a borage desaturase cDNA containing an N-terminal cytochrome b5 domain
1388 results in the accumulation of high levels of $\Delta 6$ - desaturated fatty acids in transgenic
1389 tobacco. Proc Natl Acad Sci USA 1997; 94:4211–6.

1390 [152] Aki T, Shimada Y, Inagaki K, Higashimoto H, Kawamoto S, Shiget S, et al. Molecular
1391 cloning and functional characterisation of rat $\Delta 6$ fatty acid desaturase. Biochem Biophys
1392 Res Commun 1999; 255:575-9.

1393 [153] Cho HP, Nakamura MT, Clarke SD. Cloning, expression and nutritional regulation of
1394 the human $\Delta 6$ desaturase. J Biol Chem 1999; 274:471-7.

1395 [154] Cho HP, Nakamura MT, Clarke SD. Cloning, expression and nutritional regulation of
1396 the human $\Delta 5$ desaturase. J Biol Chem 1999; 274:37335-9.

1397 [155] Zolfaghari R, Cifelli CJ, Banta MD, Ross AC. Fatty acid delta(5)-desaturase mRNA is
1398 regulated by dietary vitamin A and exogenous retinoic acid in liver of adult rats. Arch
1399 Biochem Biophys 2001; 391:8–15.

1400 [156] Matsuzaka T, Shimano H, Yahagi N, Amemiya-Kudo M, Yoshikawa T, Hasty AH, et al.
1401 Dual regulation of mouse Delta(5)- and Delta(6)-desaturase gene expression by SREBP-1
1402 and PPARalpha. J Lipid Res 2002; 43:107–14.

1403 [157] Dreesden TD, Adamson AW, Tekle M, Tang C, Cho HP, Clarke SD, Gettys TW. A
1404 newly discovered member of the fatty acid desaturase gene family: A non-coding,
1405 antisense RNA gene to $\Delta 5$ -desaturase. Prostaglandins Leukot Essent Fatty Acids 2006; 75:
1406 97-106.

1407 [158] Mathias RA, Sergeant S, Ruczinski I, Torgerson DG, Hugenschmidt CE, Kubala M, et
1408 al. The impact of FADS genetic variants on omega6 polyunsaturated fatty acid metabolism
1409 in African Americans. BMC Genet 2011; 12:50.

1410 [159] Sergeant S, Hugenschmidt CE, Rudock ME, Ziegler JT, Ivester P, Ainsworth HC, et al.
1411 Differences in arachidonic acid levels and fatty acid desaturase (FADS) gene variants in
1412 African Americans and European Americans with diabetes or the metabolic syndrome. Br
1413 J Nutr 2012; 107: 547–55.

1414 [160] Mathias RA, Fu W, Akey JM, Ainsworth HC, Torgerson DG, Ruczinski, I, et al.
1415 Adaptive evolution of the FADS gene cluster within Africa. PLoS ONE 2012; 7: e44926.

1416 [161] Oh CS, Toke DA, Mandala S, Martin CE. ELO2 and ELO3, homologues of the
 1417 *Saccharomyces cerevisiae* ELO1 gene, function in fatty acid elongation and are required
 1418 for sphingolipid formation. J Biol Chem 1997; 272:17376-84.

1419 [162] Tvrdik P, Westerberg R, Silve S, Asadi A, Jakobsson A, Cannon B, et al. Role of a new
 1420 mammalian gene family in the biosynthesis of very long chain fatty acids and
 1421 sphingolipids. J Cell Biol 2000; 149:707-18.

1422 [163] Suneja SK, Nagi MN, Cook L, Cinti DL. Decreased long-chain fatty acyl CoA
 1423 elongation activity in quaking and jimpy mouse brain: Deficiency in one enzyme or
 1424 multiple enzyme activities? J Neurochem 1991; 57:140-6.

1425 [164] Tvrdik P, Asadi A, Kozak LP, Nedergaard J, Cannon B, Jacobsson A. Cig30, a mouse
 1426 member of a novel membrane protein gene family, is involved in the recruitment of brown
 1427 adipose tissue. J Biol Chem 1997; 272:31738-46.

1428 [165] Brolinson A, Fourcade S, Jakobsson A, Pujol A, Jacobsson A. Steroid hormones control
 1429 circadian Elovl3 expression in mouse liver. Endocrinology 2008; 149:3158-66.

1430 [166] Moon YA, Shah NA, Mohapatra S, Warrington JA, Horton JD. Identification of a
 1431 mammalian long chain fatty acyl elongase regulated by sterol regulatory element-binding
 1432 proteins. J Biol Chem 2001; 276:45358-66.

1433 [167] Inagaki K, Aki T, Fukuda Y, Kawamoto S, Shigeta S, Ono K, Suzuki O. Identification
 1434 and expression of a rat fatty acid elongase involved the biosynthesis of C18 fatty acids.
 1435 Biosci Biotechnol Biochem 2002; 66:613-21.

1436 [168] Tamura K, Makino A, Hullin-Matsuda F, Kobayashi T, Furihata M, Chung S, et al.
 1437 Novel lipogenic enzyme ELOVL7 is involved in prostate cancer growth through saturated
 1438 long-chain fatty acid metabolism. Cancer Res 2009; 69:8133-40.

1439 [169] Zhang K, Kniazeva M, Han M, Li W, Yu Z, Yang Z, et al. A 5-bp deletion in ELOVL4
 1440 is associated with two related forms of autosomal dominant macular dystrophy. Nat Genet
 1441 2001; 27:89-93.

1442 [170] Zhang XM, Yang Z, Karan G, Hashimoto T, Baehr W, Yang XJ, Zhang K. Elovl4
 1443 mRNA distribution in the developing mouse retina and phylogenetic conservation of
 1444 Elovl4 genes. Mol Vis 2003; 9:301-7.

1445 [171] Cameron DJ, Tong Z, Yang Z, Kaminoh J, Kamiyah S, Chen H, Zeng J, Chen Y, Lou L,
 1446 Zhang K. Essential role of Elovl4 in very long chain fatty acid synthesis, skin permeability
 1447 barrier function, and neonatal survival. Int J Biol Sci 2007; 3:111-9.

1448 [172] Li W, Sandhoff R, Kono M, Zerfas P, Hoffman V, Ding BC-H, et al. Depletion of
1449 ceramides with very long chain fatty acids causes defective skin permeability barrier
1450 function, and neonatal lethality in ELOVL4 deficient mice. *Int J Biol Sci* 2007; 3:120-8.

1451 [173] Aveldaño MI. A novel group of very long chain polyenoic fatty acids in
1452 dipolyunsaturated phosphatidylcholines from vertebrate retina. *J. Biol. Chem.* 1987;
1453 262:1172–9.

1454 [174] Aveldaño MI. Phospholipid species containing long and very long polyenoic fatty acids
1455 remain with rhodopsin after hexane extraction of photoreceptor membranes. *Biochemistry*
1456 1988; 27:1229-39.

1457 [175] McMahon A, Jackson SN, Woods AS, Kedziński W. A Stargardt disease-3 mutation in
1458 the mouse *Elovl4* gene causes retinal deficiency of C32-C36 acyl phosphatidylcholines.
1459 *FEBS Lett* 2007; 581:5459-5463.

1460 [176] Robinson BS, Johnson DW, Poulos A. Unique molecular species of
1461 phosphatidylcholine containing very-long-chain (C24-C38) polyenoic fatty acids in rat
1462 brain. *Biochem J* 1990; 265, 763-7.

1463 [177] Aveldaño MI, Robinson BS, Johnson D, Poulos A. Long and very long chain
1464 polyunsaturated fatty acids of the n-6 series in rat seminiferous tubules. Active
1465 desaturation of 24:4n-6 to 24:5n-6 and concomitant formation of odd and even chain
1466 tetraenoic and pentaenoic fatty acids up to C32. *J Biol Chem* 1993; 268:11663–9.

1467 [178] Furland NE, Maldonado EN, Aveldaño MI. Very long chain PUFA in murine testicular
1468 triglycerides and cholesterol esters. *Lipids* 2003; 38:73-80.


1469 [179] Furland NE, Oresti GM, Antollini SS, Venturino A, Maldonado EN, Aveldaño MI.
1470 Very long-chain polyunsaturated fatty acids are the major acyl groups of sphingomyelins
1471 and ceramides in the head of mammalian spermatozoa. *J. Biol. Chem.* 2007; 282:18151-61.

1472 [180] Furland NE, Maldonado EN, Ayuza-Aresti P, Aveldaño MI. Changes in lipids
1473 containing long- and very long-chain polyunsaturated fatty acids in cryptorchid rat testes.
1474 *Biol Reprod* 2007; 77:181-8.

1475 [181] Agbaga M-P, Brush RS, Mandal MNA, Henry K, Elliott MH, Anderson RE. Role of
1476 Stargardt-3 macular dystrophy protein (ELOVL4) in the biosynthesis of very long chain
1477 fatty acids. *Proc Natl Acad Sci USA* 2008; 105:12843-8.

1478 [182] Vasireddy V, Uchida Y, Salem Jr N, Kim SY, Mandal MNA, Bhanuprakash Reddy G,
1479 et al. Loss of functional ELOVL4 depletes very long-chain fatty acids (\geq C28) and the
1480 unique ω -O-acylceramides in skin leading to neonatal death. *Hum Mol Genet*
1481 2007;16:471-82.

1482 [183] Agbaga M-P, Mandal MNA, Anderson RE. Retinal very long-chain PUFAs: new
1483 insights from studies on ELOVL4 protein. J Lipid Res 2010; 51:1624-42.
1484
1485



1487
1488 Figure 1. Fatty acid nomenclature. Examples of the A) saturated fatty acid palmitic acid or
1489 hexadecanoic acid (16:0), B) monounsaturated fatty acid oleic acid or *cis* Δ^9 -octadecenoic
1490 acid (18:1n-9), C) polyunsaturated fatty acid α -linolenic acid or *all cis* $\Delta^9,12,15$ -
1491 octadecatrienoic acid (18:3n-3), and long-chain polyunsaturated fatty acid docosahexaenoic
1492 acid or *all cis* $\Delta^4,7,10,13,16,19$ -docosahexaenoic acid (22:6n-3).
1493




Figure 2. Phylogenetic relationships of extant chordate lineages. WGD – whole genome duplications; TSWGD – teleost-specific whole genome duplications. ✓ indicates the presence of available genome sequence.




Figure 3. Fatty acid elongation pathway occurring in endoplasmic reticulum. The elongase (Elovl) is responsible for the condensation of malonyl-CoA into an activated fatty acid, regarded as the rate-limiting step in the pathway. Other enzymes involved in the pathway are 3-ketoacyl-CoA reductase (Kar), 3-hydroxyacyl-CoA dehydrase (Hadc) and *trans*-2,3,-enoyl-CoA reductase (Ter).




Figure 4. Biosynthetic pathways of polyunsaturated fatty acids. Desaturation reactions are denoted with “ ω ” or “ Δ ” to refer, respectively, the carbon position at which the incipient double bond locates within the methyl (ω) and front (Δ) ends of fatty acyl chain. Elongation reactions are catalyzed by Elovl. *De novo* biosynthesis of stearic acid (18:0) by fatty acid synthase (FAS) complex system and subsequent $\Delta 9$ unsaturation by stearoyl-CoA desaturase are also shown. Production of 22:6n-3 and 22:5n-6 occurs through partial β -oxidation of 24:6n-3 and 24:5n-6, respectively.




Figure 5. Elovl and Fads gene repertoire in extant chordate lineages. WGD – whole genome duplications; TSWGD – teleost-specific whole genome duplications. ✓ indicates the presence of full genome sequences; * indicates gene loss in various teleost species; • indicates gene loss in species examined to date; # indicates absent gene in tunicates.





Figure 6. The elephant shark *Fads* locus.



1524

1525 Figure 7. The evolution of Fads and Elovl gene repertoire in chordates. Empty boxes signal
 1526 gene loss.

1527

Tables

Table 1. Fatty acyl desaturases investigated in teleosts. The desaturation activities determined for each characterized desaturase enzyme are indicated.

Species*	Common name	Reported activities	References
<i>Danio rerio</i> ¹	Zebrafish	$\Delta 6, \Delta 5, \Delta 8$	[70, 100]
<i>Oncorhynchus mykiss</i>	Rainbow trout	$\Delta 6, \Delta 8$	[70, 99, 101]
<i>Salmo salar</i> ($\Delta 5Fad$)	Atlantic salmon	$\Delta 5$	[102]
<i>S. salar</i> ($\Delta 6Fad_a$)		$\Delta 6$	[103]
<i>S. salar</i> ($\Delta 6Fad_b$)		$\Delta 6, \Delta 8$	[70, 104]
<i>S. salar</i> ($\Delta 6Fad_c$)		$\Delta 6, \Delta 8$	[70, 104]
<i>Cyprinus carpio</i>	Common carp	$\Delta 6$	[101]
<i>Sparus aurata</i>	Gilthead sea bream	$\Delta 6, \Delta 8$	[70, 101, 105]
<i>Psetta maxima</i>	Turbot	$\Delta 6, \Delta 8$	[70, 101]
<i>Gadus morhua</i>	Atlantic cod	$\Delta 6, \Delta 8$	[70, 106]
<i>Rachycentron canadum</i>	Cobia	$\Delta 6, \Delta 8$	[70, 107]
<i>Dicentrarchus labrax</i>	European sea bass	$\Delta 6$	[108, 109]
<i>Lates calcarifer</i>	Barramundi	$\Delta 6$	[110]
<i>Siganus canaliculatus</i> (Fad1) ¹	Rabbitfish	$\Delta 6, \Delta 5, \Delta 8$	[70, 71]
<i>S. canaliculatus</i> (Fad2) ²		$\Delta 4, \Delta 5, \Delta 8$	[70, 71]
<i>Acanthopagrus schlegeli</i>	Black seabream		[111]
<i>Nibea mitsukurii</i>	Nibe croaker	$\Delta 6, \Delta 8$	[112]
<i>Thunnus thynnus</i>	Northern bluefin tuna	$\Delta 6$	[113]
<i>Solea senegalensis</i> ²	Senegalese sole	$\Delta 4, \Delta 5$	[97]
<i>Oreochromis niloticus</i> ¹	Nile tilapia	$\Delta 6, \Delta 5$	[114]
<i>Argyrosomus regius</i>	Meagre	$\Delta 6, \Delta 8$	[98]
<i>Chirostoma estor</i> (Fads2a) ¹	Pike silverside	$\Delta 4, \Delta 5$	[92]
<i>C. estor</i> (Fads2b) ²		$\Delta 6, \Delta 5, \Delta 8$	[92]
<i>Anguilla japonica</i>	Japanese eel	$\Delta 6, \Delta 8$	[115]
<i>Epinephelus coioides</i>	Orange spotted grouper	$\Delta 6, \Delta 8$	[116]
<i>Channa striata</i> ²	Striped snakehead	$\Delta 4, \Delta 5$	[117]

*In species with more than one desaturase being studied (*Salmo salar*, *Siganus canaliculatus*, *Chirostoma estor*), the name of the specific gene is indicated according to published information.

¹ Bifunctional $\Delta 6\Delta 5$ desaturases

² Bifunctional $\Delta 4\Delta 5$ desaturases

Table 2. Elongases (Elovl) investigated in teleosts. The type of Elovl is determined by amino acid similarities with mammalian orthologues and substrate specificities as reported in the indicated publications.

Species	Common name	Elongase type	Substrate specificities	References
<i>Danio rerio</i>	Zebrafish	Elovl5	C ₁₈₋₂₂	[130]
<i>D. rerio</i>		Elovl2	C ₁₈₋₂₄	[131]
<i>D. rerio</i>		Elovl4a	C ₂₀₋₃₄	[132]
<i>D. rerio</i>		Elovl4b	C ₂₀₋₃₄	[132]
<i>Salmo salar</i>	Atlantic salmon	Elovl5a	C ₁₈₋₂₂	[96, 133, 134]
<i>S. salar</i>		Elovl5b	C ₁₈₋₂₂	[133, 134]
<i>S. salar</i>		Elovl2	C ₁₈₋₂₂	[134]
<i>S. salar</i>		Elovl4b	C ₂₀₋₃₄	[135]
<i>Oreochromis niloticus</i>	Nile tilapia	Elovl5	C ₁₈₋₂₂	[136]
<i>Clarius gariepinus</i>	African catfish	Elovl5	C ₁₈₋₂₂	[133]
<i>Sparus aurata</i>	Gilthead sea bream	Elovl5	C ₁₈₋₂₂	[133]
<i>Psetta maxima</i>	Turbot	Elovl5	C ₁₈₋₂₂	[133]
<i>Gadus morhua</i>	Atlantic cod	Elovl5	C ₁₈₋₂₂	[133]
<i>Rachycentron canadum</i>	Cobia	Elovl5	C ₁₈₋₂₂	[107]
<i>R. canadum</i>		Elovl4b	C ₂₀₋₃₄	[136]
<i>Lates calcarifer</i>	Barramundi	Elovl5	C ₁₈₋₂₂	[110]
<i>Nibea mitsukurii</i>	Nibe croaker	Elovl5	C ₁₈₋₂₂	[112]
<i>N. mitsukurii</i>		Elovl4b	C ₁₈₋₂₂	[112]
<i>Thunnus maccoyii</i>	Southern bluefin tuna	Elovl5	C ₁₈₋₂₂	[137]
<i>Thunnus thynnus</i>	Northern bluefin tuna	Elovl5	C ₁₈₋₂₂	[113]
<i>Siganus canaliculatus</i>	Rabbitfish	Elovl5	C ₁₈₋₂₂	[138]
<i>S. canaliculatus</i>		Elovl4b	C ₂₀₋₃₄	[138]
<i>Solea senegalensis</i>	Senegalese sole	Elovl5	C ₁₈₋₂₂	[97]
<i>Acanthopagrus schlegelii</i>	Black seabream	Elovl5	C ₁₈₋₂₀	[139]
<i>Argyrosomus regius</i>	Meagre	Elovl5	C ₁₈₋₂₂	[98]
<i>Esox lucius</i>	Northern pike	Elovl5	C ₁₈₋₂₂	[96]
<i>Oncorhynchus mykiss</i>	Rainbow trout	Elovl5	C ₁₈₋₂₀	[140]
<i>O. mykiss</i>		Elovl2	C ₂₀₋₂₂	[140]
<i>Chirostoma estor</i>	Pike silverside	Elovl5	C ₁₈₋₂₂	[92]
<i>Anguilla japonica</i>	Japanese eel	Elovl5	C ₁₈₋₂₀	[115]

1544 Table 3. Fatty acyl desaturases (Fads) and elongases (Elovl) investigated in non-teleost
1545 chordates.

Species	Gene	Common name	References
<i>MAMMALS</i>			
<i>Homo sapiens</i>	<i>FADS1</i>	Human	[154]
<i>H. sapiens</i>	<i>FADS2</i>		[73, 127, 154]
<i>H. sapiens</i>	<i>ELOVL2</i>		[142]
<i>H. sapiens</i>	<i>ELOVL4</i>		[181]
<i>H. sapiens</i>	<i>ELOVL5</i>		[141]
<i>Mus musculus</i>	<i>Fads1</i>	Mouse	[156]
<i>M. musculus</i>	<i>Fads2</i>		[156]
<i>M. musculus</i>	<i>Elovl2</i>		[142]
<i>M. musculus</i>	<i>Elovl5</i>		[142]
<i>Rattus norvegicus</i>	<i>Fads1</i>	Rat	[155]
<i>R. norvegicus</i>	<i>Fads2</i>		[128, 152]
<i>R. norvegicus</i>	<i>Elovl2</i>		[146]
<i>R. norvegicus</i>	<i>Elovl5</i>		[146, 167]
<i>Papio anubis</i>	<i>FADS2</i>	Baboon	[69]
<i>BIRDS</i>			
<i>Gallus gallus</i>	<i>ELOVL2</i>	Chicken	[90]
<i>G. gallus</i>	<i>ELOVL5</i>		[90]
<i>Anas platyrhynchos</i>	<i>ELOVL2</i>	Domestic Mallard duck	[91]
<i>A. platyrhynchos</i>	<i>ELOVL5</i>		[91]
<i>Meleagris gallopavo</i>	<i>ELOVL2</i>	Australian white hybrid turkey	[91]
<i>M. gallopavo</i>	<i>ELOVL5</i>		[91]
<i>TUNICATES</i>			
<i>Ciona intestinalis</i>	<i>elovl4</i>	Sea squirt	[88]
<i>CHONDRICHTHYANS</i>			
<i>Scyliorhinus canicula</i>	<i>fads1</i>	Lesser-spotted dogfish	[89]
<i>S. canicula</i>	<i>fads2</i>	Lesser-spotted dogfish	[89]

1546
1547